


# Introduction to Mean-Field Theory of Spin Glass Models

Václav Janiš

Institute of Physics, Academy of Sciences of the Czech Republic, Praha, CZ

Autumn School on Correlated Electrons:  
Many-Body Physics: From Kondo to Hubbard  
21-25 September 2015, FZ Jülich



Collaborators: Anna Kauch, Antonín Klíč (FZÚ AV ČR, v. v. i.)

# Motivation

## Why mean-field theory of spin glasses?

- Diluted magnetic impurities (Mn, Fe) in a metallic matrix (Cu, Au, Ag, Pt)
- Fluctuating long-range spin exchange (RKKY) -- mean field approximation should be good
- Standard approach fails - inconsistent
- Analytic approach -- replica trick & replica-symmetry breaking (no direct physical interpretation)
- New type of long range order: non-measurable order parameters

What is the physical meaning of replica-symmetry breaking?  
Can we avoid replicas?

Fundamental concept to follow: Ergodicity



# Motivation

Why mean-field theory of spin glasses?

- Diluted magnetic impurities (Mn, Fe) in a metallic matrix (Cu, Au, Ag, Pt)
- Fluctuating long-range spin exchange (RKKY) -- mean field approximation should be good
- Standard approach fails - inconsistent
- Analytic approach -- replica trick & replica-symmetry breaking (no direct physical interpretation)
- New type of long range order: non-measurable order parameters

What is the physical meaning of replica-symmetry breaking?  
Can we avoid replicas?

Fundamental concept to follow: Ergodicity

# Motivation

Why mean-field theory of spin glasses?

- Diluted magnetic impurities (Mn, Fe) in a metallic matrix (Cu, Au, Ag, Pt)
- Fluctuating long-range spin exchange (RKKY) -- mean field approximation should be good
- Standard approach fails - inconsistent
- Analytic approach -- replica trick & replica-symmetry breaking (no direct physical interpretation)
- New type of long range order: non-measurable order parameters

What is the physical meaning of replica-symmetry breaking?  
Can we avoid replicas?

Fundamental concept to follow: Ergodicity



# Outline

- 1 Introduction - spin models and mean-field solution
  - Models of interacting spins
  - Models with disorder and frustration - spin glasses
- 2 Fundamental concepts: Ergodicity, thermodynamic homogeneity and real replicas
  - Ergodicity in statistical physics
  - Real-replica method for restoring thermodynamic homogeneity
- 3 Hierarchical construction of mean-field theory of spin glasses
  - Discrete replica-symmetry (replica-independence) breaking
  - Continuous replica-symmetry breaking
- 4 Solvable cases: 1RSB and asymptotic  $T \nearrow T_c$  solutions
  - One-level RSB -- Ising
  - Infinite RSB - asymptotic solution -- Ising
  - Potts and  $p$ -spin glass
- 5 Conclusions



# Heisenberg spins

- Model of interacting spins (Heisenberg)

$$H[J, \mathbf{S}] = - \sum_{i < j} J_{ij} \mathbf{S}_i \cdot \mathbf{S}_j$$

- Spin exchange
  - Ferromagnetic interaction:  $J_{ij} > 0$
  - Antiferromagnetic interaction:  $J_{ij} < 0$
- Regular crystalline structure (lattice)
- Strong anisotropy: Only single spin projection ( $S^z$ ) interacts
- Ising model

$$H[J, S] = - \sum_{i < j} J_{ij} S_i S_j$$

- Classical spins with  $S_i = \pm 1$  ( $\hbar/2$  units)



# Other spin models -- generalizations of Ising I

- Potts model -  $p > 2$  spin projections


$$H_p = - \sum_{i < j} J_{ij} \delta_{n_i, n_j}$$

$n_i = 1, 2, \dots, p$

- Spin representation

$$H_P [J, \mathbf{S}] = -\frac{1}{2} \sum_{i,j} J_{ij} \mathbf{S}_i \cdot \mathbf{S}_j - \sum_i \mathbf{h} \cdot \mathbf{S}_i ,$$

Potts vectors  $\mathbf{S}_i = \{s_i^1, \dots, s_i^{p-1}\}$ , values are state vectors  $\{\mathbf{e}_A\}_{A=1}^p$



# Other spin models -- generalizations of Ising II

$$\sum_{A=1}^p e_A^\alpha = 0, \quad \sum_{A=1}^p e_A^\alpha e_A^\beta = p \delta^{\alpha\beta}, \quad \sum_{\alpha=1}^{p-1} e_A^\alpha e_B^\alpha = p \delta_{AB} - 1$$

## ■ Explicit representation

$$e_A^\alpha = \begin{cases} 0 & A < \alpha \\ \sqrt{\frac{p(p-\alpha)}{p+1-\alpha}} & A = \alpha \\ \frac{1}{\alpha-p} \sqrt{\frac{p(p-\alpha)}{p+1-\alpha}} & A > \alpha . \end{cases}$$

## ■ $p$ -spin model

$$H_p [J, S] = \sum_{1 \leq i_1 < i_2 < \dots < i_p} J_{i_1 i_2 \dots i_p} S_{i_1} S_{i_2} \dots S_{i_p} .$$

$S$  are Ising spins,  $p = 2$  reduces to Ising



# Ising thermodynamics -- mean-field solution

- Thermally induced spin fluctuations -- free energy

$$-\beta F(T) = \ln \text{Tr}_S \exp \{-\beta H[J, S]\}$$

- Long-range ferromagnetic interaction:  $J_{ij} = -J/N$
- Mean-field (Weiss) solution with ergodic assumption

$$f(T, m) = F(T, m)/N = \frac{Jm^2}{2} - \frac{1}{\beta} \ln 2 \cosh(\beta Jm)$$

global magnetization  $m$  -- variational parameter

- Equilibrium state -- magnetization minimizing  $f(T, m)$
- Equilibrium magnetization

$$m = \tanh(\beta Jm)$$

minimizes free energy



# Critical point -- ergodicity & symmetry breaking



- Critical point  $\beta J = 1$  separates two phases
  - Paramagnetic:  $m = 0$
  - Ferromagnetic:  $1 \geq m^2 > 0$
- Ergodicity broken in the FM phase (in a trivial way)
- Spin-reflection symmetry  $H[J, S] = H[J, -S]$  broken
- Non-ergodic situation: degenerate solution ( $F(T, m) = F(T, -m)$ )
- Adding magnetic energy  $H'[h, S] = -h \sum_i S_i$  lifts degeneracy & restores ergodicity

Ergodicity (uniqueness of equilibrium state) restored  
by a symmetry-breaking magnetic field



# Disorder & frustration - inhomogeneous spin exchange I

- Randomness in the spin exchange
- *System locally frustrated*: ferro (red bond) and antiferro (blue bond) randomly distributed



- Unbiased situation -- neither ferro nor antiferro ordering preferred  
*Gaussian random variables* in mean-field limit

# Disorder & frustration - inhomogeneous spin exchange

II

$$N \langle J_{ij} \rangle_{av} = \sum_{j=1}^N J_{ij} = 0, \quad N \langle J_{ij}^2 \rangle_{av} = \sum_{j=1}^N J_{ij}^2 = J^2$$

## ■ Ising model

$$P(J_{ij}) = \sqrt{\frac{N}{2\pi J^2}} \exp \left\{ -\frac{NJ_{ij}^2}{2J^2} \right\}$$

## ■ Potts model (not symmetric v.r.t. spin reflection)

$$P(J_{ij}) = \sqrt{\frac{N}{2\pi J^2}} \exp \frac{-N(J_{ij} - J_0/N)^2}{2J^2},$$

$J_0 = \sum_j J_{0j}$  -- averaged (ferromagnetic) interaction



# Disorder & frustration - inhomogeneous spin exchange

III

## ■ $p$ -spin model

$$P(J_{i_1 i_2 \dots i_p}) = \sqrt{\frac{N^{p-1}}{\pi p!}} \exp \left\{ -\frac{J_{i_1 i_2 \dots i_p}^2 N^{p-1}}{J^2 p!} \right\}$$

## Real spin-glass systems

- Highly diluted magnetic ions (Fe, Mn) in noble metals (Au, Cu)
  - RKKY interaction -- generates effectively random long-range spin exchange
  - Critical behavior in magnetic field -- FC § ZFC




FIG. 7. Static susceptibilities of CuMn vs temperature for 1.08 and 2.02 at. % Mn. After zero-field cooling ( $H < 0.05$  Oe), initial susceptibilities (b) and (d) were taken for increasing temperature in a field of  $H = 5.9$  Oe. The susceptibilities (a) and (c) were obtained in the field  $H = 5.9$  Oe, which was applied above  $T_g$  before cooling the samples. From Nagata *et al.* (1979).




FIG. 1. Real part  $\Gamma'$  of the complex susceptibility  $\Gamma(\omega)$  as a function of temperature for sample IIC (CuMn with 0.94 at. % Mn, powder). Inset reveals frequency dependence and rounding of the cusp by use of strongly expanded coordinate scales. Measuring frequencies:  $\square$ , 1.33 kHz;  $\circ$ , 234 Hz;  $\triangle$ , 104 Hz;  $\times$ , 4.6 Hz. From Mulder *et al.* (1981).

# Thermodynamics of disordered and frustrated systems

## -- spin glasses

### Assumptions and basic properties of spin glass models (MFT)

- **Ergodic hypothesis** -- self averaging of thermodynamic potentials (free energy in thermodynamic limit equals the averaged one)
- **Low-temperature phase** -- local magnetic moments without homogeneous magnetic order
- **Degenerate thermodynamic state** -- ergodicity broken
- **No symmetry of the Hamiltonian broken**

How to reach thermodynamic limit (infinite volume)?  
How to restore ergodicity? What are the order parameters?

# Thermodynamics of disordered and frustrated systems

## -- spin glasses

### Assumptions and basic properties of spin glass models (MFT)

- **Ergodic hypothesis** -- self averaging of thermodynamic potentials (free energy in thermodynamic limit equals the averaged one)
- **Low-temperature phase** -- local magnetic moments without homogeneous magnetic order
- **Degenerate thermodynamic state** -- ergodicity broken
- **No symmetry of the Hamiltonian broken**

How to reach thermodynamic limit (infinite volume)?  
How to restore ergodicity? What are the order parameters?



# Sherrington-Kirkpatrick mean-field solution

- Averaged free energy -- replica trick with ergodic assumption (apart from critical point)

$$\beta F = -\langle \ln Z \rangle_{av} = -\lim_{\nu \rightarrow 0} \left[ \frac{1}{\nu} \lim_{N \rightarrow \infty} (\langle Z_N^\nu \rangle_{av} - 1) \right].$$

- Single order parameter  $q = N^{-1} \sum_i m_i^2$
- Mean-field replica-symmetric solution: free-energy density

$$f(T, q) = -\frac{\beta}{4}(1 - q)^2 - \frac{1}{\beta} \int_{-\infty}^{\infty} D\eta \ln 2 \cosh [\beta(h + \eta\sqrt{q})]$$

- Global parameter:  $q = N^{-1} \sum_i m_i^2 = \langle \tanh^2 [\beta(h + \eta\sqrt{q})] \rangle_\eta$   
maximizes free energy!

- Inconsistency:

- Zero temperature entropy negative:  $S(0) = -\sqrt{\frac{2}{\pi}} k_B \approx -0.798 k_B$
- Instability in the low-temperature phase:

$$\Lambda = 1 - \beta^2 \left\langle (1 - \tanh^2 [\beta(h + \eta\sqrt{q})])^2 \right\rangle_\eta < 0$$



# Sherrington-Kirkpatrick mean-field solution

- Averaged free energy -- replica trick with ergodic assumption (apart from critical point)

$$\beta F = -\langle \ln Z \rangle_{av} = -\lim_{\nu \rightarrow 0} \left[ \frac{1}{\nu} \lim_{N \rightarrow \infty} (\langle Z_N^\nu \rangle_{av} - 1) \right].$$

- Single order parameter  $q = N^{-1} \sum_i m_i^2$
- Mean-field replica-symmetric solution: free-energy density

$$f(T, q) = -\frac{\beta}{4}(1 - q)^2 - \frac{1}{\beta} \int_{-\infty}^{\infty} d\eta \ln 2 \cosh [\beta(h + \eta\sqrt{q})]$$

- Global parameter:  $q = N^{-1} \sum_i m_i^2 = \langle \tanh^2 [\beta(h + \eta\sqrt{q})] \rangle_\eta$   
maximizes free energy!
- Inconsistency:

- Zero temperature entropy negative:  $S(0) = -\sqrt{\frac{2}{\pi}} k_B \approx -0.798 k_B$
- Instability in the low-temperature phase:

$$\Lambda = 1 - \beta^2 \left\langle (1 - \tanh^2 [\beta(h + \eta\sqrt{q})])^2 \right\rangle_\eta < 0$$



## 1 Introduction - spin models and mean-field solution

- Models of interacting spins
- Models with disorder and frustration - spin glasses

## 2 Fundamental concepts: Ergodicity, thermodynamic homogeneity and real replicas

- Ergodicity in statistical physics
- Real-replica method for restoring thermodynamic homogeneity

## 3 Hierarchical construction of mean-field theory of spin glasses

- Discrete replica-symmetry (replica-independence) breaking
- Continuous replica-symmetry breaking

## 4 Solvable cases: 1RSB and asymptotic $T \nearrow T_c$ solutions

- One-level RSB -- Ising
- Infinite RSB - asymptotic solution -- Ising
- Potts and  $p$ -spin glass

## 5 Conclusions



# Ergodicity in equilibrium statistical physics

## ■ Fundamental ergodic theorem (Birkhoff)

$$\langle f \rangle_T \equiv \lim_{T \rightarrow \infty} \frac{1}{T} \int_{t_0}^{t_0+T} f(X(t)) dt = \frac{1}{\Sigma_E} \int_{S_E} f(X) dS_E \equiv \langle f \rangle_S$$

## ■ Phase space homogeneously covered by the phase trajectory

$$\lim_{T \rightarrow \infty} \frac{\tau_R}{T} = \frac{\Sigma_R(E)}{\Sigma(E)}$$

## ■ Equilibrium ergodic macroscopic state

- homogeneously spread over the allowed phase space
- characterized by homogeneous parameters ( $\{E, T\}$ ,  $\{N, \mu\}$ , ...)
- number of relevant parameters (Legendre pairs)  $\text{\'a priori}$  unknown

How do we determine the phase space  
covered by the phase space trajectory?



# Homogeneity of thermodynamic potentials

## ■ Homogeneity in the phase space

$$S(E) = k_B \ln \Gamma(E) = \frac{k_B}{\nu} \ln \Gamma(E)^\nu = \frac{k_B}{\nu} \ln \Gamma(\nu E)$$

$$F(T) = -\frac{k_B T}{\nu} \ln [\text{Tr } e^{-\beta H}]^\nu = -\frac{k_B T}{\nu} \ln [\text{Tr } e^{-\beta \nu H}]$$

## ■ Homogeneity of thermodynamic potentials (Euler)

$$\alpha F(T, V, N, \dots, X_i, \dots) = F(T, \alpha V, \alpha N, \dots, \alpha X_i, \dots)$$

Density of the free energy  $f = F/N$

-- function of only **densities** of extensive variables  $X_i/N$

Ergodicity (homogeneity) guarantees existence and uniqueness of the thermodynamic limit  $N \rightarrow \infty$



# Homogeneity of thermodynamic potentials

## ■ Homogeneity in the phase space

$$S(E) = k_B \ln \Gamma(E) = \frac{k_B}{\nu} \ln \Gamma(E)^\nu = \frac{k_B}{\nu} \ln \Gamma(\nu E)$$

$$F(T) = -\frac{k_B T}{\nu} \ln [\text{Tr } e^{-\beta H}]^\nu = -\frac{k_B T}{\nu} \ln [\text{Tr } e^{-\beta \nu H}]$$

## ■ Homogeneity of thermodynamic potentials (Euler)

$$\alpha F(T, V, N, \dots, X_i, \dots) = F(T, \alpha V, \alpha N, \dots, \alpha X_i, \dots)$$

Density of the free energy  $f = F/N$

-- function of only **densities** of extensive variables  $X_i/N$

Ergodicity (homogeneity) guarantees existence and uniqueness of the thermodynamic limit  $N \rightarrow \infty$



# Ergodicity breaking

- Ergodicity gives meaning to statistical averages
- Thermodynamic properties in the infinite-volume limit
- Ergodicity breaking -- improper statistical phase space
  - 1 caused by a phase transition breaking a symmetry of the Hamiltonian
  - 2 without apparent symmetry breaking -- glass-like behavior
- Means to restore ergodicity
  - 1 Measurable (physical) symmetry breaking fields
  - 2 Real replicas (non-measurable symmetry breaking fields)

Ergodicity must be restored to establish stable equilibrium



# Ergodicity breaking

- Ergodicity gives meaning to statistical averages
- Thermodynamic properties in the infinite-volume limit
- Ergodicity breaking -- improper statistical phase space
  - 1 caused by a phase transition breaking a symmetry of the Hamiltonian
  - 2 without apparent symmetry breaking -- glass-like behavior
- Means to restore ergodicity
  - 1 Measurable (physical) symmetry breaking fields
  - 2 Real replicas (non-measurable symmetry breaking fields)

Ergodicity must be restored to establish stable equilibrium



# Real replicas -- stability w.r.t. phase-space scalings

Real replicas -- means to probe thermodynamic homogeneity

Replicated Hamiltonian:  $[H]_\nu = \sum_{a=1}^{\nu} H^a = \sum_{\alpha=1}^{\nu} \sum_{<ij>} J_{ij} S_i^a S_j^a$

Symmetry-breaking fields:  $\Delta H(\mu) = \frac{1}{2} \sum_{a \neq b} \sum_i \mu^{ab} S_i^a S_i^b$

Averaged replicated free energy with coupled replicas

$$F_\nu(\mu) = -k_B T \frac{1}{\nu} \left\langle \ln \text{Tr} \exp \left\{ -\beta \sum_a H^a - \beta \Delta H(\mu) \right\} \right\rangle_{av}$$

Analytic continuation to non-integer parameter  $\nu$

Stability w.r.t. phase space scaling:  $\lim_{\mu \rightarrow 0} \frac{dF_\nu(\mu)}{d\nu} \equiv 0$

Real replicas - simulate impact of surrounding bath



# Real replicas -- stability w.r.t. phase-space scalings

Real replicas -- means to probe thermodynamic homogeneity

Replicated Hamiltonian:  $[H]_\nu = \sum_{a=1}^{\nu} H^a = \sum_{\alpha=1}^{\nu} \sum_{<ij>} J_{ij} S_i^a S_j^a$

Symmetry-breaking fields:  $\Delta H(\mu) = \frac{1}{2} \sum_{a \neq b} \sum_i \mu^{ab} S_i^a S_i^b$

Averaged replicated free energy with coupled replicas

$$F_\nu(\mu) = -k_B T \frac{1}{\nu} \left\langle \ln \text{Tr} \exp \left\{ -\beta \sum_a H^a - \beta \Delta H(\mu) \right\} \right\rangle_{av}$$

Analytic continuation to non-integer parameter  $\nu$

Stability w.r.t. phase space scaling:

$$\lim_{\mu \rightarrow 0} \frac{dF_\nu(\mu)}{d\nu} \equiv 0$$

Real replicas - simulate impact of surrounding bath



# Annealed vs. quenched disorder

- Averaged ( $\nu$ -times replicated) partition function

$$\langle Z'_N \rangle_{av} = \int D[J] \mu[J] \prod_{a=1}^{\nu} \prod_{i=1}^N d[\mathbf{S}_i^a] \rho[\mathbf{S}_i^a] \exp \left\{ -\beta \sum_{a=1}^{\nu} H[J, \mathbf{S}^a] \right\}$$

- Averaged ( $\nu$ -times replicated) free energy

$$-\beta \langle F'_N \rangle_{av} = \int D[J] \mu[J] \ln \int \prod_{a=1}^{\nu} \prod_{i=1}^N d[\mathbf{S}_i^a] \rho[\mathbf{S}_i^a] \exp \left\{ -\beta \sum_{a=1}^{\nu} H[J, \mathbf{S}^a] \right\}$$

- Replicas for disordered systems:

- Quenched disorder (spin glasses) -- replica trick ( $\nu \rightarrow 0$ )

$$\beta F_{qu} = - \lim_{\nu \rightarrow 0} \left[ \frac{1}{\nu} \lim_{N \rightarrow \infty} (\langle Z'_N \rangle_{av} - 1) \right]$$

- Annealed disorder -- thermodynamic homogeneity ( $\nu$  arbitrary)

$$\beta F_{an} = - \frac{1}{\nu} \lim_{N \rightarrow \infty} \ln \langle Z'_N \rangle_{av}$$



- 1** Introduction - spin models and mean-field solution
  - Models of interacting spins
  - Models with disorder and frustration - spin glasses
- 2** Fundamental concepts: Ergodicity, thermodynamic homogeneity and real replicas
  - Ergodicity in statistical physics
  - Real-replica method for restoring thermodynamic homogeneity
- 3** Hierarchical construction of mean-field theory of spin glasses
  - Discrete replica-symmetry (replica-independence) breaking
  - Continuous replica-symmetry breaking
- 4** Solvable cases: 1RSB and asymptotic  $T \nearrow T_c$  solutions
  - One-level RSB -- Ising
  - Infinite RSB - asymptotic solution -- Ising
  - Potts and  $p$ -spin glass
- 5** Conclusions



# Ergodicity breaking -- broken LRT in replicated space

- Breaking of LRT to inter-replica interaction  $\mu^{ab} \rightarrow 0$

$$f_\nu = \frac{\beta \mathcal{P}}{4} \left[ \frac{1}{\nu} \sum_{a \neq b}^{\nu} \left\{ (\chi^{ab})^2 + 2q\chi^{ab} \right\} - (1-q)^2 \right] - \frac{1}{\beta\nu} \int_{-\infty}^{\infty} \frac{d\eta}{\sqrt{2\pi}} e^{-\eta^2/2} \ln \text{Tr}_\nu \exp \left\{ \beta^2 \mathcal{P} \sum_{a < b}^{\nu} \chi^{ab} S^a S^b + \beta \bar{h} \sum_{a=1}^{\nu} S^a \right\}$$

$$\chi^{ab} = \langle \langle S^a S^b \rangle_T \rangle_{av} - q, \quad q = \langle \langle S^a \rangle_T^2 \rangle_{av}, \quad \bar{h} = h + \eta \sqrt{q}$$

- Free energy  $f_\nu$  must be analytic function of index  $\nu$
- Parisi conditions for analytic continuation

$$\chi^{aa} = 0, \quad \chi^{ab} = \chi^{ba}, \quad \sum_{c=1}^{\nu} (\chi^{ac} - \chi^{bc}) = 0$$

- $K < \nu - 1$  different inter-replica susceptibilities  $\chi_1, \dots, \chi_K$  with multiplicities  $\nu_1, \dots, \nu_K$

https://www.mathworksheetsland.com/grade-3/area/area-rectangle.html



# Ergodicity breaking -- broken LRT in replicated space

- Breaking of LRT to inter-replica interaction  $\mu^{ab} \rightarrow 0$

$$f_\nu = \frac{\beta J^2}{4} \left[ \frac{1}{\nu} \sum_{a \neq b}^{\nu} \left\{ (\chi^{ab})^2 + 2q\chi^{ab} \right\} - (1-q)^2 \right] - \frac{1}{\beta\nu} \int_{-\infty}^{\infty} \frac{d\eta}{\sqrt{2\pi}} e^{-\eta^2/2} \ln \text{Tr}_\nu \exp \left\{ \beta^2 J^2 \sum_{a < b}^{\nu} \chi^{ab} S^a S^b + \beta \bar{h} \sum_{a=1}^{\nu} S^a \right\}$$

$$\chi^{ab} = \langle \langle S^a S^b \rangle_T \rangle_{av} - q, \quad q = \langle \langle S^a \rangle_T^2 \rangle_{av}, \quad \bar{h} = h + \eta \sqrt{q}$$

- Free energy  $f_\nu$  must be analytic function of index  $\nu$
- Parisi conditions for analytic continuation

$$\chi^{aa} = 0, \quad \chi^{ab} = \chi^{ba}, \quad \sum_{c=1}^{\nu} (\chi^{ac} - \chi^{bc}) = 0$$

- $K < \nu - 1$  different inter-replica susceptibilities  $\chi_1, \dots, \chi_K$  with multiplicities  $\nu_1, \dots, \nu_K$



# Analytic continuation

- Only specific matrices  $\nu \times \nu$  allow for analytic continuation to real  $\nu$
- Multiplicity of the order parameters –  $K$  different values

$$\begin{pmatrix} 0 & q_0 & 0 & 0 \\ q_0 & 0 & q_0 & q_0 \\ 0 & q_0 & 0 & q_0 \\ q_0 & q_0 & q_0 & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$q = q_0 + q_1, \quad \eta = 2^{1/2} (q - 1) = 2^{1/2} q$$



# Analytic continuation

- Only specific matrices  $\nu \times \nu$  allow for analytic continuation to real  $\nu$
- Multiplicity of the order parameters --  $K$  different values

$$\begin{pmatrix} 0 & q_0 & q_1 & q_1 & q_2 & q_2 & q_2 & q_2 \\ q_0 & 0 & q_1 & q_1 & q_2 & q_2 & q_2 & q_2 \\ q_1 & q_1 & 0 & q_0 & q_2 & q_2 & q_2 & q_2 \\ q_1 & q_1 & q_0 & 0 & q_2 & q_2 & q_2 & q_2 \\ q_2 & q_2 & q_2 & q_2 & 0 & q_0 & q_1 & q_1 \\ q_2 & q_2 & q_2 & q_2 & q_0 & 0 & q_1 & q_1 \\ q_2 & q_2 & q_2 & q_2 & q_1 & q_1 & 0 & q_0 \\ q_2 & q_2 & q_2 & q_2 & q_1 & q_1 & q_0 & 0 \end{pmatrix}$$

$$q_I = q + \chi_I, \nu_I = 2^I, \nu - 1 = \sum_I^K \nu_I$$

# Analytic continuation

- Only specific matrices  $\nu \times \nu$  allow for analytic continuation to real  $\nu$
- Multiplicity of the order parameters --  $K$  different values

$$\begin{pmatrix} 0 & q_0 & q_1 & q_1 & q_2 & q_2 & q_2 & q_2 \\ q_0 & 0 & q_1 & q_1 & q_2 & q_2 & q_2 & q_2 \\ q_1 & q_1 & 0 & q_0 & q_2 & q_2 & q_2 & q_2 \\ q_1 & q_1 & q_0 & 0 & q_2 & q_2 & q_2 & q_2 \\ q_2 & q_2 & q_2 & q_2 & 0 & q_0 & q_1 & q_1 \\ q_2 & q_2 & q_2 & q_2 & q_0 & 0 & q_1 & q_1 \\ q_2 & q_2 & q_2 & q_2 & q_1 & q_1 & 0 & q_0 \\ q_2 & q_2 & q_2 & q_2 & q_1 & q_1 & q_0 & 0 \end{pmatrix}$$

$$q_I = q + \chi_I, \nu_I = 2^I, \nu - 1 = \sum_I^K \nu_I$$



# Analytic continuation

- Only specific matrices  $\nu \times \nu$  allow for analytic continuation to real  $\nu$
- Multiplicity of the order parameters --  $K$  different values

$$\begin{pmatrix} 0 & q_0 & q_1 & q_1 & q_2 & q_2 & q_2 & q_2 \\ q_0 & 0 & q_1 & q_1 & q_2 & q_2 & q_2 & q_2 \\ q_1 & q_1 & 0 & q_0 & q_2 & q_2 & q_2 & q_2 \\ q_1 & q_1 & q_0 & 0 & q_2 & q_2 & q_2 & q_2 \\ q_2 & q_2 & q_2 & q_2 & 0 & q_0 & q_1 & q_1 \\ q_2 & q_2 & q_2 & q_2 & q_0 & 0 & q_1 & q_1 \\ q_2 & q_2 & q_2 & q_2 & q_1 & q_1 & 0 & q_0 \\ q_2 & q_2 & q_2 & q_2 & q_1 & q_1 & q_0 & 0 \end{pmatrix}$$

$$q_I = q + \chi_I, \nu_I = 2^I, \nu - 1 = \sum_I^K \nu_I$$



# Analytic continuation

- Only specific matrices  $\nu \times \nu$  allow for analytic continuation to real  $\nu$
- Multiplicity of the order parameters --  $K$  different values

$$\begin{pmatrix} 0 & q_0 & q_1 & q_1 & q_2 & q_2 & q_2 & q_2 \\ q_0 & 0 & q_1 & q_1 & q_2 & q_2 & q_2 & q_2 \\ q_1 & q_1 & 0 & q_0 & q_2 & q_2 & q_2 & q_2 \\ q_1 & q_1 & q_0 & 0 & q_2 & q_2 & q_2 & q_2 \\ q_2 & q_2 & q_2 & q_2 & 0 & q_0 & q_1 & q_1 \\ q_2 & q_2 & q_2 & q_2 & q_0 & 0 & q_1 & q_1 \\ q_2 & q_2 & q_2 & q_2 & q_1 & q_1 & 0 & q_0 \\ q_2 & q_2 & q_2 & q_2 & q_1 & q_1 & q_0 & 0 \end{pmatrix}$$

$$q_I = q + \chi_I, \nu_I = 2^I, \nu - 1 = \sum_I^K \nu_I$$



# Analytic continuation

- Only specific matrices  $\nu \times \nu$  allow for analytic continuation to real  $\nu$
- Multiplicity of the order parameters --  $K$  different values


$$\begin{pmatrix} 0 & q_0 & q_1 & q_1 & q_2 & q_2 & q_2 & q_2 \\ q_0 & 0 & q_1 & q_1 & q_2 & q_2 & q_2 & q_2 \\ q_1 & q_1 & 0 & q_0 & q_2 & q_2 & q_2 & q_2 \\ q_1 & q_1 & q_0 & 0 & q_2 & q_2 & q_2 & q_2 \\ q_2 & q_2 & q_2 & q_2 & 0 & q_0 & q_1 & q_1 \\ q_2 & q_2 & q_2 & q_2 & q_0 & 0 & q_1 & q_1 \\ q_2 & q_2 & q_2 & q_2 & q_1 & q_1 & 0 & q_0 \\ q_2 & q_2 & q_2 & q_2 & q_1 & q_1 & q_0 & 0 \end{pmatrix}$$

$$q_I = q + \chi_I, \nu_I = 2^I, \nu - 1 = \sum_I^K \nu_I$$

# ultrametric structure

## ultrametric structure

- only block matrices of identical elements
- larger blocks multiples of smaller blocks
- hierarchy of embeddings around diagonal
- ultrametric metrics (tree-like)



# Multiple replica hierarchies

Averaged free energy density with  $K$  hierarchies of replicas

$$\Delta\chi_I = \chi_I - \chi_{I+1} \geq \Delta\chi_{I+1} \geq 0, \quad \nu_I \text{ -- arbitrary positive}$$

$$f_K(q; \Delta\chi_1, \dots, \Delta\chi_K, \nu_1, \dots, \nu_K) = -\frac{\beta}{4} \left(1 - q - \sum_{I=1}^K \Delta\chi_I\right)^2 - \frac{1}{\beta} \ln 2 \\ + \frac{\beta}{4} \sum_{I=1}^K \nu_I \Delta\chi_I \left[2 \left(q + \sum_{i=I}^K \Delta\chi_i\right) - \Delta\chi_I\right] - \frac{1}{\beta} \int_{-\infty}^{\infty} \mathcal{D}\eta \ln Z_K$$


Hierarchical local partition sums  $Z_I = \left[\int_{-\infty}^{\infty} \mathcal{D}\lambda_I Z_{I-1}^{\nu_I}\right]^{1/\nu_I}$

Initial condition  $Z_0 = \cosh \left[ \beta \left( h + \eta \sqrt{q} + \sum_{I=1}^K \lambda_I \sqrt{\Delta\chi_I} \right) \right]$

Gaussian measure  $\mathcal{D}\lambda \equiv d\lambda e^{-\lambda^2/2}/\sqrt{2\pi}$



# Multiple embeddings -- including boundary terms



- $\Delta\chi_I$  -- inter-replica interaction strength,
- $\lambda_I$  -- effective magnetic field due to replicated spins
- $\nu_I V$ : volume affected by replicated spins -- range of inter-replica interaction

$$\frac{N}{V} \ln Z_{I-1}(\beta, \bar{h}_I)$$

$$\rightarrow \frac{N}{\nu_I V} \ln \int \mathcal{D}\lambda_I Z_{I-1}^{\nu_I} (\beta, \bar{h}_I + \lambda_I \sqrt{\Delta\chi_I})$$

- Effective weight of surrounding spins in thermal averaging

$$\rho_I = \frac{Z_{I-1}^{\nu_I}}{\langle Z_{I-1}^{\nu_I} \rangle_{\lambda_I}}$$

# Equilibrium state -- stationarity equations & stability

- Stationarity equations with discrete  $K$  replica hierarchies

$$q = \langle \langle t \rangle_K^2 \rangle_\eta ,$$

$$\Delta\chi_I = \langle \langle \langle t \rangle_{I-1}^2 \rangle_K \rangle_\eta - \langle \langle \langle t \rangle_I^2 \rangle_K \rangle_\eta ,$$

$$\nu_I \Delta\chi_I = \frac{4}{\beta^2} \frac{\langle \langle \ln Z_{I-1} \rangle_K \rangle_\eta - \langle \langle \ln Z_I \rangle_K \rangle_\eta}{2 \left( q + \sum_{i=I+1}^K \Delta\chi_i \right) + \Delta\chi_I}$$

$$t \equiv \tanh \left[ \beta \left( h + \eta \sqrt{q} + \sum_{I=1}^K \lambda_I \sqrt{\Delta\chi_I} \right) \right] ,$$

$$\langle t \rangle_I(\eta; \lambda_K, \dots, \lambda_{I+1}) = \langle \rho_I \dots \langle \rho_1 t \rangle_{\lambda_1} \dots \rangle_{\lambda_I}, \quad \rho_I = Z_{I-1}^{\nu_I} / \langle Z_{I-1}^{\nu_I} \rangle_{\lambda_I}$$

- $K+1$  stability conditions determine number  $K$

$$\Lambda_I^K = 1 - \beta^2 \left\langle \left\langle \left\langle 1 - t^2 + \sum_{i=0}^I \nu_i (\langle t \rangle_{i-1}^2 - \langle t \rangle_i^2) \right\rangle_I \right\rangle_K \right\rangle_\eta \geq 0$$



# Infinite many replica hierarchies I

Limit to infinite number of replica hierarchies  $K \rightarrow \infty$

- Infinitesimal differences  $\Delta\chi_I$  and  $\Delta\nu_I$ :

$$\Delta\chi_I = \Delta\chi/K, \Delta\nu_I = \Delta m/K, \Delta\chi_I/\Delta\nu_I \rightarrow x(m) < \infty$$

- Parisi continuous free energy (around 1RSB):

$$f(q, \chi_1, m_1, m_0; x(m)) = -\frac{\beta}{4}(1-q-\chi_1-X_0(m_1))^2 + \frac{\beta}{4} \left[ m_1 (q + \chi_1 + X_0(m_1))^2 - m_0 q^2 \right] - \frac{\beta}{4} \int_{m_0}^{m_1} dm [q + \chi_1 + X_0(m)]^2 - \frac{1}{\beta} \langle g_1(m_0, h + \eta\sqrt{q}) \rangle_\eta$$

- Integral representation of the interacting part

$$g_1(m_0, h) = \mathbb{E}_0(m_0, m_1, h) \circ g_1(h)$$

$$\equiv \overline{\mathbb{T}}_m \exp \left\{ \frac{1}{2} \int_{m_0}^{m_1} dm x(m) [\partial_{\bar{h}}^2 + mg'_1(m; h + \bar{h}) \partial_{\bar{h}}] \right\} g_1(h + \bar{h}) \Big|_{\bar{h}=0}$$

$$g'_1(m; h) = \partial g_1(m, h) / \partial h$$



# Infinite many replica hierarchies II

- Anti time-ordering product from quantum many-body PT

$$\overline{\mathbb{T}}_\lambda \exp \left\{ \int_0^1 d\lambda \widehat{O}(\lambda) \right\} \equiv 1 + \sum_{n=1}^{\infty} \int_0^1 d\lambda_1 \int_0^{\lambda_1} \dots \int_0^{\lambda_{n-1}} d\lambda_n \widehat{O}(\lambda_n) \dots \widehat{O}(\lambda_1)$$

- Initial condition (1RSB)

$$g_1(h) \equiv g_1(m_1, h) = \frac{1}{m_1} \ln \int_{-\infty}^{\infty} \frac{d\phi}{\sqrt{2\pi}} e^{-\phi^2/2} [2 \cosh (\beta(h + \phi\sqrt{\chi_1}))]^{m_1}$$

- Closed implicit equation

$$g'_1(m, h) = \mathbb{E}(m, m_1, h) \circ g_0(h)$$

$$\equiv \overline{\mathbb{T}}_m \exp \left\{ \frac{1}{2} \int_m^{m_1} dm' x(m') [\partial_{\bar{h}}^2 + 2m' g'_1(m'; h + \bar{h}) \partial_{\bar{h}}] \right\} g'_1(h + \bar{h}) \Big|_{\bar{h}=0}$$

- Notation:  $X_0(m) = \int_{m_0}^m dm' x(m')$

using the notation of the previous slide, we have  $\mathbb{E}(m, m_1, h) = X_0(m) + \int_m^{m_1} dm' x(m') g'_1(m'; h + \bar{h})$



# Discrete vs. continuous replica-symmetry breaking

## Discrete RSB

- 1 Hierarchical embeddings -- ultrametric structure
- 2 No restriction on the replica-induced order parameters
- 3 The number of replica hierarchies  $K$  from stability conditions
- 4 Either unstable or locally stable

## Continuous RSB

- 1 Limit of infinite number of replica hierarchies
- 2 Infinitesimal distance between replica hierarchies
- 3 Closed theory independently of stability of the discrete scheme
- 4 Always marginally stable



- 1 Introduction - spin models and mean-field solution
  - Models of interacting spins
  - Models with disorder and frustration - spin glasses
- 2 Fundamental concepts: Ergodicity, thermodynamic homogeneity and real replicas
  - Ergodicity in statistical physics
  - Real-replica method for restoring thermodynamic homogeneity
- 3 Hierarchical construction of mean-field theory of spin glasses
  - Discrete replica-symmetry (replica-independence) breaking
  - Continuous replica-symmetry breaking
- 4 Solvable cases: 1RSB and asymptotic  $T \nearrow T_c$  solutions
  - One-level RSB -- Ising
  - Infinite RSB - asymptotic solution -- Ising
  - Potts and  $p$ -spin glass
- 5 Conclusions



# First level replica-symmetry (ergodicity) breaking

- Ergodicity broken in the SG phase -- one embedding

$$f(q; \chi, \nu) = -\frac{\beta}{4}(1-q)^2 + \frac{\beta}{4}(\nu-1)\chi(2q+\chi) + \frac{\beta}{2}\chi - \frac{1}{\beta\nu} \int_{-\infty}^{\infty} D\eta \ln \int_{-\infty}^{\infty} D\lambda \{2 \cosh [\beta(h + \eta\sqrt{q} + \lambda\sqrt{\chi})]\}^{\nu}$$

- stationarity equations ( $t \equiv \tanh[\beta(h + \eta\sqrt{q} + \lambda\sqrt{\chi})]$ )

$$\begin{aligned} q &= \langle \langle t \rangle_{\lambda}^2 \rangle_{\eta}, & q_{EA} &= q + \chi = \langle \langle t^2 \rangle_{\lambda} \rangle_{\eta} \\ \beta^2 \chi (2q + \chi) \nu &= [\langle \ln \cosh[\beta(h + \eta\sqrt{q} + \lambda\sqrt{\chi})] \rangle_{\lambda} \\ &\quad - \ln \langle \cosh^{\nu}[\beta(h + \eta\sqrt{q} + \lambda\sqrt{\chi})] \rangle_{\lambda}^{1/\nu}] \end{aligned}$$


- Stability conditions

$$\Lambda_0 = 1 - \beta^2 \left\langle \left\langle (1-t)^2 \right\rangle_{\lambda} \right\rangle_{\eta}$$


$$\Lambda_1 = 1 - \beta^2 \left\langle \left\langle 1 - (1-\nu)t^2 - \nu \langle t^2 \rangle_{\lambda} \right\rangle_{\lambda}^2 \right\rangle_{\eta}$$




# 1RSB - thermodynamics



# RS & 1RSB - instability



Instability at zero magnetic field



SG phase in magnetic field - AT line

# SK model at zero magnetic field

Only asymptotic expansions available for  $K \rightarrow \infty$

- Small expansion parameter  $\tau = (T_c - T)/T_c$

$$\Delta\chi_I^K \doteq \frac{2}{2K+1} \tau, \quad \nu_I^K \doteq \frac{4(K-I+1)}{2K+1} \tau, \quad q^K \doteq \frac{1}{2K+1} \tau,$$

$$Q^K \equiv q_{EA} = q + \chi_1 - \chi_K \doteq \tau + \frac{12K(K+1)+1}{3(2K+1)^2} \tau^2, \quad \Lambda_I^K \doteq -\frac{4}{3} \frac{\tau^2}{(2K+1)^2}$$

$$\chi_T \doteq \beta \left( 1 - Q^K + \sum_{I=1}^K m_I \Delta\chi_I \right) \doteq 1 - \frac{\tau^2}{3(2K+1)^2}$$

$$\Delta f \doteq \left( \frac{1}{6}\tau^3 + \frac{7}{24}\tau^4 + \frac{29}{120}\tau^5 \right) - \frac{1}{360}\tau^5 \left( \frac{1}{K} \right)^4$$

Parisi continuous ansatz proven right

# SK model in magnetic field

- Full RSB at AT line reduces to 1RSB ( $h > 0$ )
- Small expansion parameter  $\alpha = \beta^2 \langle (1 - t_0^2)^2 \rangle_\eta - 1$   
( $t_0 \equiv \tanh[\beta(h + \eta\sqrt{q})]$ )

$$\nu = \frac{2\langle t_0^2(1 - t_0^2)^2 \rangle_\eta}{\langle (1 - t_0^2)^3 \rangle_\eta}$$

$$\chi_1 = \frac{1}{2\beta^2\nu} \frac{\beta^2 \langle (1 - t_0^2)^2 \rangle_\eta - 1}{1 - 3\beta^2 \langle t_0^2(1 - t_0^2)^2 \rangle_\eta} + O(\alpha^2)$$

$$\nu_I^K \doteq \nu_1 + (K+1-2I)\Delta\nu/K, \quad \Delta\chi_I^K \doteq \chi_1/K$$

$$\Delta\nu \doteq \frac{\beta^2\chi_1 \left\langle \left(1 - t_0^2\right)^2 \left(2\left(1 - 3t_0^2\right)^2 + 3\left(t_0^2 - 1\right)\nu\left(8t_0^2 + \left(t_0^2 - 1\right)\nu\right)\right) \right\rangle_\eta}{\langle (1 - t_0^2)^3 \rangle_\eta}$$

$$\Lambda_I^K \doteq -\frac{2\beta^2}{3K^2} \frac{\chi_1 \Delta\nu}{\nu + 2}$$



# Potts glass ( $p < 4$ ): discrete RSB

- Two 1RSB solutions for  $\nu_1 \doteq \frac{p-2}{2} + \frac{36-12p+p^2}{8(4-p)}\tau$

- Locally stable solution (near  $T_c$  and  $p > p^* \approx 2.82$ )

$$q^{(1)} \doteq 0, \quad \Delta\chi^{(1)} \doteq \frac{2}{4-p}\tau$$

stability function:  $\Lambda_1^{(1)} \doteq \frac{\tau^2(p-1)}{6(4-p)^2} (7p^2 - 24p + 12)$

- Unstable solution ( $p > p^*$  unphysical)

$$q^{(2)} \doteq \frac{-12 + 24p - 7p^2}{3(4-p)^2(p-2)}\tau^2, \quad \Delta\chi^{(2)} \doteq \frac{2}{4-p}\tau$$

- $K_{RSB}$  (from the unstable one)


$$q^K \doteq -\frac{1}{3K^2} \frac{12 - 24p + 7p^2}{(4-p)^2(p-2)}\tau^2, \quad \Delta\chi_I^K \doteq \frac{1}{K} \frac{2}{(4-p)}\tau,$$

$$\nu_I^K \doteq \frac{p-2}{2} + \frac{2}{4-p} \left[ 3 + \frac{3}{2}p - p^2 + \left( 3 - 6p + \frac{7}{4}p^2 \right) \frac{2I-1}{2K} \right] \tau$$



# Potts glass ( $p = 3$ ): coexistence


1RSB & FRSB coexist near  $T_c$



Stability and entropy of 1RSB solution ( $p = 3$ )

Free-energy differences:

$$\beta(f_c - f_{1RSB}) \doteq \frac{(p-1)(p(7p-24)+12)^2\tau^5}{720(4-p)^5}, \quad \beta(f_c - f_{RS}) \doteq \frac{(p-1)(p-2)^2\tau^3}{3(4-p)(6-p)^2}$$



Free-energy difference as function of  $p$



# $p$ -spin glass: 1RSB 1

- Discontinuous transition to the low-temperature phase for  $p > 2$
- Asymptotic solution  $p \rightarrow \infty$ : 1RSB

$$f_T^{(p \rightarrow \infty)}(q, \chi_1, \mu_1)$$

$$\begin{aligned} &= -\frac{1}{4T} [1 - (q + \chi_1)(1 - \ln(q + \chi_1))] - \frac{1}{\mu_1} \ln[2 \cosh(\mu_1 h)] \\ &\quad - \frac{\mu_1}{4} [\chi_1 - (q + \chi_1) \ln(q + \chi_1)] - \frac{\mu_1 q}{4} [\ln q + p(1 - \tanh^2(\mu_1 h))] \end{aligned}$$

rescaled variable  $\mu_1 = \beta \nu_1$

- Low-temperature solution ( $p = \infty$ ) -- Random energy model

$$\chi_1 = 1 - q, \quad q = \exp\{-p(1 - \tanh^2(\mu_1 h))\},$$


$$\mu_1 = 2\sqrt{\ln[2 \cosh(\mu_1 h)] - h \tanh(\mu_1 h)}$$

for  $\beta > 2\sqrt{\ln[2 \cosh(\beta h)] - h \tanh(\beta h)}$ ,

otherwise  $q + \chi_1 = 0$  and  $\mu_1 = \beta$

# $p$ -spin glass: 1RSB II

- Negative entropy for  $p < \infty$ 
  - full continuous free energy around 1RSB needed



# Conclusions

Spin-glass phase: Ergodicity breaking without symmetry breaking

- 1 Frustration with disorder prevents existence of physical symmetry-breaking fields
- 2 Real replicas -- means to test thermodynamic homogeneity (ergodicity)
- 3 Analytic continuation to non-integer replication index mandatory -- ultrametric structure
- 4 Broken LRT of inter-replica interaction -- broken replica symmetry (ergodicity)
- 5 Hierarchical replications -- series of admissible solutions (equilibrium states)
- 6 Local and global stability conditions select the true equilibrium
- 7 Continuous RSB -- marginally stable  
*(available only via asymptotic expansions)*

