


# Exchange Mechanisms

Erik Koch

Institute for Advanced Simulation, Forschungszentrum Jülich



# Magnetism is Quantum Mechanical

---

QUANTUM MECHANICS

THE KEY TO UNDERSTANDING MAGNETISM

Nobel Lecture, 8 December, 1977

J.H. VAN VLECK

Harvard University, Cambridge, Massachusetts, USA

## **Bohr – van Leeuwen theorem**

in a classical system in thermal equilibrium  
a magnetic field will not induce a magnetic moment

Lorentz force perpendicular to velocity  $\Rightarrow$  does not change kinetic energy

Boltzmann statistics occupies states according to energy

# magnetic moments


---

complex wave function: current density

$$\vec{j}(\vec{r}) = -\frac{e\hbar}{2im_e} \left( \overline{\psi(\vec{r})} \nabla \psi(\vec{r}) - \psi(\vec{r}) \nabla \overline{\psi(\vec{r})} \right)$$

orbital magnetic moment

$$\vec{\mu} = \frac{1}{2} \int \vec{r} \times \vec{j} d^3 = -\frac{e\hbar}{2m_e} \langle \vec{L} \rangle = -\mu_B \langle \vec{L} \rangle$$



electron spin


$$\vec{\mu}_S = -g_e \mu_B \langle \vec{S} \rangle, \quad g_e \approx 2.0023 \dots$$

**atomic moments of the order of  $\mu_B$**

# magnetic interaction


dipole-dipole interaction

$$\Delta E = \frac{\vec{\mu}_1 \cdot \vec{\mu}_2 - 3(\hat{R} \cdot \vec{\mu}_1)(\hat{R} \cdot \vec{\mu}_2)}{4\pi\epsilon_0 c^2 R^3}$$



interaction energy of two dipoles  $\mu_B$  two Bohr radii  $a_0$  apart:

$$\Delta E = -\frac{2\mu_B^2}{4\pi\epsilon_0 c^2 (2a_0)^3} = -\frac{1/2}{137^2 8} \text{ Hartree} \approx 0.09 \text{ meV}$$



**expect magnetic ordering below temperatures of about 1 K**


what about magnetite ( $\text{Fe}_3\text{O}_4$ )  
with  $T_c \approx 840 \text{ K}$  ?



# exchange mechanisms

---

coupling of magnetic moments  
results from the interplay of  
the **Pauli principle**  
with **Coulomb repulsion**  
and **electron hopping**



not a fundamental but an **effective interaction**: model/mechanism

# Models and Mechanisms

---

The art of model-building is the exclusion of real but irrelevant parts of the problem, and entails hazards for the builder and the reader. The builder may leave out something genuinely relevant; the reader, armed with too sophisticated an experimental probe or too accurate a computation, may take literally a schematized model whose main aim is to be a demonstration of possibility.



P.W. Anderson  
Local Moments and Localized States  
Nobel Lecture 1977

# Coulomb Exchange

---

Coulomb repulsion between electrons



$$H_U = \sum_{i < j} \frac{1}{|\vec{r}_i - \vec{r}_j|}$$

consider two electrons in orthogonal orbitals  $\phi_a$  and  $\phi_b$

Slater determinant of spin-orbitals:

$$\Psi_{a,\sigma; b\sigma'}(\vec{r}_1, s_1; \vec{r}_2, s_2) = \frac{1}{\sqrt{2}} \begin{vmatrix} \phi_a(\vec{r}_1) \sigma(s_1) & \phi_a(\vec{r}_2) \sigma(s_2) \\ \phi_b(\vec{r}_1) \sigma'(s_1) & \phi_b(\vec{r}_2) \sigma'(s_2) \end{vmatrix}$$

$$= \frac{1}{\sqrt{2}} \left( \phi_a(\vec{r}_1) \phi_a(\vec{r}_2) \sigma(s_1) \sigma'(s_2) - \phi_b(\vec{r}_1) \phi_a(\vec{r}_2) \sigma'(s_1) \sigma(s_2) \right)$$



# Coulomb exchange: same spin



when electrons have same spin:  $\sigma = \sigma'$

$$\psi_{a,\sigma; b\sigma} = \frac{1}{\sqrt{2}} \left( \phi_a(\vec{r}_1) \phi_b(\vec{r}_2) - \phi_b(\vec{r}_1) \phi_a(\vec{r}_2) \right) \sigma(s_1) \sigma(s_2)$$

Coulomb matrix-element

$$\left\langle \psi_{a,\sigma; b,\sigma} \left| \frac{1}{|\vec{r}_1 - \vec{r}_2|} \right| \psi_{a,\sigma; b,\sigma} \right\rangle = \frac{1}{2} (U_{ab} - J_{ab} - J_{ba} + U_{ba}) = U_{ab} - J_{ab}$$

Coulomb integral

$$U_{ab} = \int d^3 r_1 \int d^3 r_2 \frac{|\phi_a(\vec{r}_1)|^2 |\phi_b(\vec{r}_2)|^2}{|\vec{r}_1 - \vec{r}_2|}$$

exchange integral

$$J_{ab} = \int d^3 r_1 \int d^3 r_2 \frac{\overline{\phi_a(\vec{r}_1)} \phi_b(\vec{r}_1) \overline{\phi_b(\vec{r}_2)} \phi_a(\vec{r}_2)}{|\vec{r}_1 - \vec{r}_2|}$$



# Coulomb exchange: opposite spin



when electrons have opposite spin:  $\sigma = -\sigma'$

$$\psi_{a,\uparrow;b\downarrow}(\vec{r}_1, s_1; \vec{r}_2, s_2) = \frac{1}{\sqrt{2}} \left( \phi_a(\vec{r}_1)\phi_b(\vec{r}_2) \uparrow(s_1)\downarrow(s_2) - \phi_b(\vec{r}_1)\phi_a(\vec{r}_2) \downarrow(s_1)\uparrow(s_2) \right)$$

$$\psi_{a,\downarrow;b\uparrow}(\vec{r}_1, s_1; \vec{r}_2, s_2) = \frac{1}{\sqrt{2}} \left( \phi_a(\vec{r}_1)\phi_b(\vec{r}_2) \downarrow(s_1)\uparrow(s_2) - \phi_b(\vec{r}_1)\phi_a(\vec{r}_2) \uparrow(s_1)\downarrow(s_2) \right)$$

diagonal matrix-elements  $\left\langle \psi_{a,\sigma;b,-\sigma} \left| \frac{1}{|\vec{r}_1 - \vec{r}_2|} \right| \psi_{a,\sigma;b,-\sigma} \right\rangle = U_{ab}$

off-diagonal matrix-elements  $\left\langle \psi_{a\uparrow;b\downarrow} \left| \frac{1}{|\vec{r}_1 - \vec{r}_2|} \right| \psi_{a\downarrow;b\uparrow} \right\rangle = -J_{ab}$

Coulomb matrix

$$\begin{pmatrix} U_{ab} & -J_{ab} \\ -J_{ab} & U_{ab} \end{pmatrix}$$

# Coulomb exchange

$$H_U = \begin{pmatrix} U_{ab} - J_{ab} & 0 & 0 & 0 \\ 0 & U_{ab} & -J_{ab} & 0 \\ 0 & -J_{ab} & U_{ab} & 0 \\ 0 & 0 & 0 & U_{ab} - J_{ab} \end{pmatrix} \begin{matrix} \uparrow\uparrow \\ \uparrow\downarrow \\ \downarrow\uparrow \\ \downarrow\downarrow \end{matrix}$$

eigenstates

**triplet:**  $\Delta\varepsilon_{triplet} = U_{ab} - J_{ab}$

$$\psi_{\uparrow\uparrow} = \frac{1}{\sqrt{2}} \left( \phi_a(\vec{r}_1)\phi_b(\vec{r}_2) - \phi_b(\vec{r}_1)\phi_a(\vec{r}_2) \right) \quad |\uparrow\uparrow\rangle$$

$$\frac{1}{\sqrt{2}} \left( \psi_{\uparrow\downarrow} + \psi_{\downarrow\uparrow} \right) = \frac{1}{\sqrt{2}} \left( \phi_a(\vec{r}_1)\phi_b(\vec{r}_2) - \phi_b(\vec{r}_1)\phi_a(\vec{r}_2) \right) \frac{1}{\sqrt{2}} \left( |\downarrow\uparrow\rangle + |\uparrow\downarrow\rangle \right)$$

$$\psi_{\downarrow\downarrow} = \frac{1}{\sqrt{2}} \left( \phi_a(\vec{r}_1)\phi_b(\vec{r}_2) - \phi_b(\vec{r}_1)\phi_a(\vec{r}_2) \right) \quad |\downarrow\downarrow\rangle$$

**singlet:**  $\Delta\varepsilon_{singlet} = U_{ab} + J_{ab}$


$$\frac{1}{\sqrt{2}} \left( \psi_{\uparrow\downarrow} - \psi_{\downarrow\uparrow} \right) = \frac{1}{\sqrt{2}} \left( \phi_a(\vec{r}_1)\phi_b(\vec{r}_2) + \phi_b(\vec{r}_1)\phi_a(\vec{r}_2) \right) \frac{1}{\sqrt{2}} \left( |\downarrow\uparrow\rangle - |\uparrow\downarrow\rangle \right)$$

# Coulomb exchange


orthogonal orbitals  $\varphi_a$  and  $\varphi_b$ :  $J_{ab} > 0$

singlet

triplet



first of **Hund's rules**: ground-state has maximum spin




more electrons  
more complicated  
Coulomb matrix

**Multiplets in  
Transition Metal Ions**

# atomic multiplets

Q. Zhang:  
 Calculations of Atomic Multiplets  
 across the Periodic Table  
 MSc thesis, RWTH Aachen 2014  
[www.cond-mat.de/sims/multiplet](http://www.cond-mat.de/sims/multiplet)

## Multiplet calculation




$^5D$

$$E = F^{(0)} [15] + F^{(2)} \left[-\frac{5}{7}\right] + F^{(4)} \left[-\frac{5}{7}\right]$$

- $|2, 2, 2, 2\rangle = c_{21}^{\uparrow} c_{-21}^{\downarrow} c_{-11}^{\uparrow} c_{01}^{\downarrow} c_{11}^{\uparrow} c_{21}^{\downarrow} |0\rangle$
- $|2, 2, 2, 1\rangle = \frac{1}{\sqrt{4}} (c_{11}^{\uparrow} c_{21}^{\downarrow} c_{-21}^{\uparrow} c_{-11}^{\downarrow} c_{01}^{\uparrow} c_{21}^{\downarrow} - c_{01}^{\uparrow} c_{21}^{\downarrow} c_{-21}^{\uparrow} c_{-11}^{\downarrow} c_{11}^{\uparrow} c_{21}^{\downarrow} + c_{-11}^{\uparrow} c_{21}^{\downarrow} c_{-21}^{\uparrow} c_{01}^{\downarrow} c_{11}^{\uparrow} c_{21}^{\downarrow} - c_{-21}^{\uparrow} c_{21}^{\downarrow} c_{-11}^{\uparrow} c_{01}^{\downarrow} c_{11}^{\uparrow} c_{21}^{\downarrow})$
- $|2, 2, 2, 0\rangle = \frac{1}{\sqrt{6}} (c_{01}^{\uparrow} c_{11}^{\downarrow} c_{21}^{\uparrow} c_{-21}^{\downarrow} c_{-11}^{\uparrow} c_{21}^{\downarrow} - c_{-11}^{\uparrow} c_{11}^{\downarrow} c_{21}^{\uparrow} c_{-21}^{\downarrow} c_{01}^{\uparrow} c_{21}^{\downarrow} + c_{-21}^{\uparrow} c_{11}^{\downarrow} c_{21}^{\uparrow} c_{-11}^{\downarrow} c_{01}^{\uparrow} c_{21}^{\downarrow} + c_{-11}^{\uparrow} c_{01}^{\downarrow} c_{21}^{\uparrow} c_{-21}^{\downarrow} c_{11}^{\uparrow} c_{21}^{\downarrow})$
- $|2, 2, 2, -1\rangle = \frac{1}{\sqrt{4}} (c_{-11}^{\uparrow} c_{01}^{\downarrow} c_{11}^{\uparrow} c_{21}^{\downarrow} c_{-21}^{\uparrow} c_{21}^{\downarrow} - c_{-21}^{\uparrow} c_{01}^{\downarrow} c_{11}^{\uparrow} c_{21}^{\downarrow} c_{-11}^{\uparrow} c_{21}^{\downarrow} + c_{-21}^{\uparrow} c_{-11}^{\downarrow} c_{11}^{\uparrow} c_{21}^{\downarrow} c_{01}^{\uparrow} c_{21}^{\downarrow} - c_{-21}^{\uparrow} c_{-11}^{\downarrow} c_{01}^{\downarrow} c_{21}^{\uparrow} c_{11}^{\uparrow} c_{21}^{\downarrow})$
- $|2, 2, 2, -2\rangle = c_{-21}^{\uparrow} c_{-11}^{\downarrow} c_{01}^{\uparrow} c_{11}^{\downarrow} c_{21}^{\uparrow} c_{21}^{\downarrow} |0\rangle$
- $|2, 1, 2, 2\rangle = c_{11}^{\uparrow} c_{-21}^{\downarrow} c_{-11}^{\uparrow} c_{01}^{\downarrow} c_{11}^{\uparrow} c_{21}^{\downarrow} |0\rangle$
- $|2, 1, 2, 1\rangle = \frac{1}{\sqrt{4}} (c_{11}^{\uparrow} c_{21}^{\downarrow} c_{-21}^{\uparrow} c_{-11}^{\downarrow} c_{01}^{\uparrow} c_{11}^{\downarrow} - c_{01}^{\uparrow} c_{11}^{\downarrow} c_{-21}^{\uparrow} c_{-11}^{\downarrow} c_{11}^{\uparrow} c_{21}^{\downarrow} + c_{-11}^{\uparrow} c_{11}^{\downarrow} c_{-21}^{\uparrow} c_{01}^{\downarrow} c_{11}^{\uparrow} c_{21}^{\downarrow} - c_{-21}^{\uparrow} c_{11}^{\downarrow} c_{-11}^{\uparrow} c_{01}^{\downarrow} c_{11}^{\uparrow} c_{21}^{\downarrow})$
- $|2, 1, 2, 0\rangle = \frac{1}{\sqrt{6}} (c_{01}^{\uparrow} c_{11}^{\downarrow} c_{21}^{\uparrow} c_{-21}^{\downarrow} c_{-11}^{\uparrow} c_{11}^{\downarrow} - c_{-11}^{\uparrow} c_{11}^{\downarrow} c_{21}^{\uparrow} c_{-21}^{\downarrow} c_{01}^{\uparrow} c_{11}^{\downarrow} + c_{-21}^{\uparrow} c_{11}^{\downarrow} c_{21}^{\uparrow} c_{-11}^{\downarrow} c_{01}^{\uparrow} c_{11}^{\downarrow} + c_{-11}^{\uparrow} c_{01}^{\downarrow} c_{11}^{\downarrow} c_{-21}^{\uparrow} c_{11}^{\uparrow} c_{21}^{\downarrow})$
- $|2, 1, 2, -1\rangle = \frac{1}{\sqrt{4}} (c_{-11}^{\uparrow} c_{01}^{\downarrow} c_{11}^{\uparrow} c_{21}^{\downarrow} c_{-21}^{\uparrow} c_{11}^{\downarrow} - c_{-21}^{\uparrow} c_{01}^{\downarrow} c_{11}^{\uparrow} c_{21}^{\downarrow} c_{-11}^{\uparrow} c_{11}^{\downarrow} + c_{-21}^{\uparrow} c_{-11}^{\downarrow} c_{11}^{\uparrow} c_{21}^{\downarrow} c_{01}^{\uparrow} c_{11}^{\downarrow} - c_{-21}^{\uparrow} c_{-11}^{\downarrow} c_{01}^{\downarrow} c_{21}^{\uparrow} c_{11}^{\uparrow} c_{21}^{\downarrow})$
- $|2, 1, 2, -2\rangle = c_{-21}^{\uparrow} c_{-11}^{\downarrow} c_{01}^{\uparrow} c_{11}^{\downarrow} c_{21}^{\uparrow} c_{11}^{\downarrow} |0\rangle$



## Self-consistent field computation



# kinetic exchange


**Coulomb** exchange: Coulomb **matrix** for anti-symmetric wave functions

**kinetic** exchange: only diagonal  **$U$** , interplay of Pauli principle and **hopping**

toy model — two sites with a single orbital

hopping between orbitals:  $t$

two electrons in same orbital:  $U$




one electron Hamiltonian (tight-binding)

$$H = \begin{pmatrix} 0 & -t \\ -t & 0 \end{pmatrix} \begin{array}{l} |\uparrow, \cdot\rangle \\ |\cdot, \uparrow\rangle \end{array}$$

eigenstates

$$\phi_{\pm} = \frac{1}{\sqrt{2}} \begin{pmatrix} \phi_1 \pm \phi_2 \end{pmatrix} \quad \varepsilon_{\pm} = \mp t$$

# direct exchange: same spin



two electrons of same spin: basis states  $|\uparrow, \uparrow\rangle, |\downarrow, \downarrow\rangle$

Hamiltonian: no hopping, no Coulomb matrix element (Pauli principle)

$$H = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \begin{array}{l} |\uparrow, \uparrow\rangle \\ |\downarrow, \downarrow\rangle \end{array}$$

$$\epsilon_{\text{triplet}} = 0$$

# direct exchange: opposite spin



two electrons of opposite spin: basis states

$|\uparrow, \downarrow\rangle, |\downarrow, \uparrow\rangle$  (covalent states)       $|\uparrow\downarrow, \cdot\rangle, |\cdot, \uparrow\downarrow\rangle$  (ionic states)

Hamiltonian

$$H = \begin{pmatrix} 0 & 0 & -t & -t \\ 0 & 0 & +t & +t \\ -t & +t & U & 0 \\ -t & +t & 0 & U \end{pmatrix} \begin{array}{l} |\uparrow, \downarrow\rangle \\ |\downarrow, \uparrow\rangle \\ |\uparrow\downarrow, \cdot\rangle \\ |\cdot, \uparrow\downarrow\rangle \end{array}$$

hopping  $-t$ : keep track of **Fermi sign!**

$$|\uparrow, \downarrow\rangle \xrightarrow{-t} |\uparrow\downarrow, \cdot\rangle \quad |\downarrow, \uparrow\rangle \xrightarrow{-(-t)} |\uparrow\downarrow, \cdot\rangle$$

# direct exchange: opposite spin




eigenstates

$$\epsilon_{\pm} = \frac{U}{2} \pm \frac{\sqrt{U^2 + 16t^2}}{2}, \quad \psi_{\pm} = \frac{\left( |\uparrow, \downarrow\rangle - |\downarrow, \uparrow\rangle - \frac{\epsilon_{\pm}}{2t} [ |\uparrow\downarrow, \cdot\rangle + |\cdot, \uparrow\downarrow\rangle ] \right)}{\sqrt{2 + \epsilon_{\pm}^2/(2t^2)}}$$

$$\epsilon_{\text{cov}} = 0, \quad \psi_{\text{cov}} = \frac{1}{\sqrt{2}} \left( |\uparrow, \downarrow\rangle + |\downarrow, \uparrow\rangle \right) \quad (\epsilon_{\text{triplet}})$$

$$\epsilon_{\text{ion}} = U, \quad \psi_{\text{ion}} = \frac{1}{\sqrt{2}} \left( |\uparrow\downarrow, \cdot\rangle - |\cdot, \uparrow\downarrow\rangle \right)$$



limit  $U \rightarrow \infty$  (or  $t \rightarrow 0$ ):

$$\epsilon_{-} \rightarrow U + 4t^2/U$$

$$\epsilon_{+} \rightarrow -4t^2/U$$



# downfolding

partition Hilbert space

$$H = \begin{pmatrix} H_{00} & T_{01} \\ T_{10} & H_{11} \end{pmatrix}$$




resolvent

$$G(\varepsilon) = (\varepsilon - H)^{-1} = \begin{pmatrix} \varepsilon - H_{00} & -T_{01} \\ -T_{10} & \varepsilon - H_{11} \end{pmatrix}^{-1}$$



inverse of 2x2 block-matrix

$$G_{00}(\varepsilon) = \left( \varepsilon - \left[ H_{00} + T_{01}(\varepsilon - H_{11})^{-1}T_{10} \right] \right)^{-1}$$



downfolded Hamiltonian

$$H_{\text{eff}} \approx H_{00} + T_{01}(\varepsilon_0 - H_{11})^{-1}T_{10}$$

good approximation: narrow energy range and/or small coupling

# inversion by partitioning

---

2×2 matrix

$$M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \quad M^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$$

invert block-2×2 matrix

solve

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \quad M^{-1} = \begin{pmatrix} \tilde{A} & \tilde{B} \\ \tilde{C} & \tilde{D} \end{pmatrix} \quad \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} \tilde{A} & \tilde{B} \\ \tilde{C} & \tilde{D} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$A\tilde{A} + B\tilde{C} = 1 \quad = (A - BD^{-1}C)\tilde{A}$$

$$C\tilde{A} + D\tilde{C} = 0 \rightsquigarrow \tilde{C} = -D^{-1}C\tilde{A}$$

$$\rightsquigarrow \tilde{A} = (A - BD^{-1}C)^{-1}$$

# direct exchange: effective Hamiltonian

systematic treatment of limit  $U \rightarrow \infty$  (or  $t \rightarrow 0$ ): downfolding

$$H = \left( \begin{array}{cc|cc} 0 & 0 & -t & -t \\ 0 & 0 & +t & +t \\ \hline -t & +t & U & 0 \\ -t & +t & 0 & U \end{array} \right)$$

downfolding eliminates ionic states (actually change of basis)


$$H_{\text{eff}}(\varepsilon) = \begin{pmatrix} -t & -t \\ +t & +t \end{pmatrix} \begin{pmatrix} \varepsilon - U & 0 \\ 0 & \varepsilon - U \end{pmatrix}^{-1} \begin{pmatrix} -t & +t \\ -t & +t \end{pmatrix} \approx -\frac{2t^2}{U} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}$$

diagonalize  $H_{\text{eff}}$

$$\begin{array}{lll} \varepsilon_t = 0 & \psi_t = \frac{1}{\sqrt{2}} \left( |\uparrow, \downarrow\rangle + |\downarrow, \uparrow\rangle \right) & \text{triplet} \\ \varepsilon_s = -\frac{4t^2}{U} & \psi_s = \frac{1}{\sqrt{2}} \left( |\uparrow, \downarrow\rangle - |\downarrow, \uparrow\rangle \right) & \text{singlet} \end{array}$$

# direct exchange: effective spin-coupling

$$J_{\text{direct}} = \varepsilon_{\text{triplet}} - \varepsilon_{\text{singlet}} = 4t^2/U \quad J > 0 \quad \text{AF coupling}$$



effective spin-Hamiltonian

$$H_{\text{eff}} = -\frac{2t^2}{U} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \begin{array}{l} |\uparrow, \downarrow\rangle \\ |\downarrow, \uparrow\rangle \end{array}$$

$$= +\frac{2t^2}{U} \left( 2S_1^z S_2^z - \frac{1}{2} + (S_1^+ S_2^- + S_1^- S_2^+) \right) = \frac{4t^2}{U} \left( \vec{S}_1 \cdot \vec{S}_2 - \frac{1}{4} \right)$$

Heisenberg  $J$

# keeping track of all these signs...

---

## towards second quantization

Slater determinant  $\Phi_{\alpha\beta}(x_1, x_2) = \frac{1}{\sqrt{2}} (\varphi_{\alpha}(x_1)\varphi_{\beta}(x_2) - \varphi_{\beta}(x_1)\varphi_{\alpha}(x_2))$

corresponding Dirac state  $|\alpha, \beta\rangle = \frac{1}{\sqrt{2}} (|\alpha\rangle|\beta\rangle - |\beta\rangle|\alpha\rangle)$

use operators  $|\alpha, \beta\rangle = c_{\beta}^{\dagger} c_{\alpha}^{\dagger} |0\rangle$

position of operators encodes signs

$$c_{\beta}^{\dagger} c_{\alpha}^{\dagger} |0\rangle = |\alpha, \beta\rangle = -|\beta, \alpha\rangle = -c_{\alpha}^{\dagger} c_{\beta}^{\dagger} |0\rangle$$

product of operators changes sign when commuted: anti-commutation

anti-commutator  $\{A, B\} := AB + BA$

# second quantization: motivation

---

specify  $N$ -electron states using operators

$N=0$ :  $|0\rangle$  (vacuum state)

normalization:  $\langle 0|0\rangle = 1$

$N=1$ :  $|\alpha\rangle = c_\alpha^\dagger|0\rangle$  (creation operator adds one electron)

normalization:  $\langle \alpha|\alpha\rangle = \langle 0|c_\alpha c_\alpha^\dagger|0\rangle$

overlap:  $\langle \alpha|\beta\rangle = \langle 0|c_\alpha c_\beta^\dagger|0\rangle$

adjoint of creation operator removes one electron:  
annihilation operator

$$c_\alpha|0\rangle = 0 \text{ and } c_\alpha c_\beta^\dagger = \pm c_\beta^\dagger c_\alpha + \langle \alpha|\beta\rangle$$

$N=2$ :  $|\alpha, \beta\rangle = c_\beta^\dagger c_\alpha^\dagger|0\rangle$

antisymmetry:  $c_\alpha^\dagger c_\beta^\dagger = -c_\beta^\dagger c_\alpha^\dagger$

# second quantization: formalism

vacuum state  $|0\rangle$

and

set of operators  $c_\alpha$  related to single-electron states  $\varphi_\alpha(x)$

defined by:

$$c_\alpha |0\rangle = 0 \quad \{c_\alpha, c_\beta\} = 0 = \{c_\alpha^\dagger, c_\beta^\dagger\}$$

$$\langle 0|0\rangle = 1 \quad \{c_\alpha, c_\beta^\dagger\} = \langle \alpha|\beta\rangle$$


creators/annihilators operate in Fock space  
transform like orbitals!

field operators

$$\hat{\psi}(x) = \sum_n \varphi_{\alpha_n}(x) c_{\alpha_n}$$

Slater determinant

$$\frac{1}{\sqrt{N!}} \langle 0 | \hat{\psi}(x_1) \hat{\psi}(x_2) \dots \hat{\psi}(x_N) c_{\alpha_N}^\dagger \dots c_{\alpha_2}^\dagger c_{\alpha_1}^\dagger | 0 \rangle$$



# second quantization: examples

---

two-site model with one electron

$$H = -t \left( c_{1\uparrow}^\dagger c_{2\uparrow} + c_{2\uparrow}^\dagger c_{1\uparrow} + c_{1\downarrow}^\dagger c_{2\downarrow} + c_{2\downarrow}^\dagger c_{1\downarrow} \right) = -t \sum_{i,j,\sigma} c_{j\sigma}^\dagger c_{i\sigma}$$

two-site model with two electrons

$$\begin{aligned} H &= -t \left( c_{1\uparrow}^\dagger c_{2\uparrow} + c_{2\uparrow}^\dagger c_{1\uparrow} + c_{1\downarrow}^\dagger c_{2\downarrow} + c_{2\downarrow}^\dagger c_{1\downarrow} \right) + U \left( n_{1\uparrow} n_{1\downarrow} + n_{2\uparrow} n_{2\downarrow} \right) \\ &= -t \sum_{i,j,\sigma} c_{j\sigma}^\dagger c_{i\sigma} + U \sum_i n_{i\uparrow} n_{i\downarrow} \end{aligned}$$

also works for single electron and for more sites

- easy to handle Slater determinants
- easy to write down many-body Hamiltonian  
(become independent of particle number)



# Hartree-Fock

---

ansatz: Slater determinant

$$|\Psi(\theta_{\uparrow}, \theta_{\downarrow})\rangle = \left( \sin(\theta_{\downarrow}) c_{1\downarrow}^{\dagger} + \cos(\theta_{\downarrow}) c_{2\downarrow}^{\dagger} \right) \left( \sin(\theta_{\uparrow}) c_{1\uparrow}^{\dagger} + \cos(\theta_{\uparrow}) c_{2\uparrow}^{\dagger} \right) |0\rangle$$

energy expectation value

$$E(\theta_{\uparrow}, \theta_{\downarrow}) = -2t (\sin \theta_{\uparrow} \sin \theta_{\downarrow} + \cos \theta_{\uparrow} \cos \theta_{\downarrow}) (\cos \theta_{\uparrow} \sin \theta_{\downarrow} + \sin \theta_{\uparrow} \cos \theta_{\downarrow}) \\ + U (\sin^2 \theta_{\uparrow} \sin^2 \theta_{\downarrow} + \cos^2 \theta_{\uparrow} \cos^2 \theta_{\downarrow})$$

minimize wrt  $\theta_{\uparrow}$  and  $\theta_{\downarrow}$

HF orbitals respect symmetry of model: restricted Hartree-Fock (RHF)


$$\text{here: } \theta_{\uparrow} = \theta_{\downarrow} = \pi/4$$

HF allowed to break symmetry: unrestricted Hartree-Fock (UHF)

$$\text{here: } \theta_{\downarrow} = \pi/2 - \theta_{\uparrow}$$




# Hartree-Fock




# direct kinetic exchange

---

singlet




triplet



direct exchange


virtual hopping  $-t^2/U \times 2$

# superexchange



symmetry:  
only one oxygen- $p$   
involved in hopping

TMOs: negligible direct hopping  
between  $d$ -orbitals  
instead hopping via oxygen



$$H = \sum_{\sigma} \left( \varepsilon_d \sum_i n_{i\sigma} + \varepsilon_p n_{p\sigma} - t_{pd} \sum_i \left( c_{i\sigma}^{\dagger} c_{p\sigma} + c_{p\sigma}^{\dagger} c_{i\sigma} \right) \right) + U_d \sum_i n_{i\uparrow} n_{i\downarrow}$$

# superexchange: same spin



$$H = \sum_{\sigma} \left( \varepsilon_d \sum_i n_{i\sigma} + \varepsilon_p n_{p\sigma} - t_{pd} \sum_i \left( c_{i\sigma}^{\dagger} c_{p\sigma} + c_{p\sigma}^{\dagger} c_{i\sigma} \right) \right) + U_d \sum_i n_{i\uparrow} n_{i\downarrow}$$

oxygen- $p$  full, two  $d$ -electrons of same spin


$$H = \left( \begin{array}{c|cc} 0 & t_{pd} & t_{pd} \\ \hline t_{pd} & U_d + \Delta_{pd} & 0 \\ t_{pd} & 0 & U_d + \Delta_{pd} \end{array} \right) \begin{array}{l} c_{2\uparrow}^{\dagger} c_{p\downarrow}^{\dagger} c_{p\uparrow}^{\dagger} c_{1\uparrow}^{\dagger} |0\rangle \\ c_{2\uparrow}^{\dagger} c_{p\uparrow}^{\dagger} c_{1\downarrow}^{\dagger} c_{1\uparrow}^{\dagger} |0\rangle \\ c_{2\downarrow}^{\dagger} c_{2\uparrow}^{\dagger} c_{p\uparrow}^{\dagger} c_{1\uparrow}^{\dagger} |0\rangle \end{array} \begin{array}{ccc} \uparrow & \uparrow\downarrow & \uparrow \\ \uparrow\downarrow & \uparrow & \uparrow \end{array}$$

$$H_{\text{eff}} = (t_{pd}, t_{pd}) \begin{pmatrix} \varepsilon - (U_d + \Delta_{pd}) & 0 \\ 0 & \varepsilon - (U_d + \Delta_{pd}) \end{pmatrix} \begin{pmatrix} t_{pd} \\ t_{pd} \end{pmatrix} \approx -\frac{2t_{pd}^2}{U_d + \Delta_{pd}}$$


# superexchange: opposite spin




0	0	$+t_{pd}$	$+t_{pd}$	0	0	0	0	0	$c_{2\downarrow}^\dagger c_{p\downarrow}^\dagger c_{p\uparrow}^\dagger c_{1\uparrow}^\dagger  0\rangle$
0	0	0	0	$+t_{pd}$	$+t_{pd}$	0	0	0	$c_{2\uparrow}^\dagger c_{p\downarrow}^\dagger c_{p\uparrow}^\dagger c_{1\downarrow}^\dagger  0\rangle$
$+t_{pd}$	0	$U_d + \Delta_{pd}$	0	0	0	$-t_{pd}$	0	$-t_{pd}$	$c_{2\downarrow}^\dagger c_{p\uparrow}^\dagger c_{1\downarrow}^\dagger c_{1\uparrow}^\dagger  0\rangle$
$+t_{pd}$	0	0	$U_d + \Delta_{pd}$	0	0	0	$-t_{pd}$	$-t_{pd}$	$c_{2\downarrow}^\dagger c_{2\uparrow}^\dagger c_{p\downarrow}^\dagger c_{1\uparrow}^\dagger  0\rangle$
0	$+t_{pd}$	0	0	$U_d + \Delta_{pd}$	0	$+t_{pd}$	0	$+t_{pd}$	$c_{2\uparrow}^\dagger c_{p\downarrow}^\dagger c_{1\downarrow}^\dagger c_{1\uparrow}^\dagger  0\rangle$
0	$+t_{pd}$	0	0	0	$U_d + \Delta_{pd}$	0	$+t_{pd}$	$+t_{pd}$	$c_{2\downarrow}^\dagger c_{2\uparrow}^\dagger c_{p\uparrow}^\dagger c_{1\downarrow}^\dagger  0\rangle$
0	0	$-t_{pd}$	0	$+t_{pd}$	0	$U_d$	0	0	$c_{p\downarrow}^\dagger c_{p\uparrow}^\dagger c_{1\downarrow}^\dagger c_{1\uparrow}^\dagger  0\rangle$
0	0	0	$-t_{pd}$	0	$+t_{pd}$	0	$U_d$	0	$c_{2\downarrow}^\dagger c_{2\uparrow}^\dagger c_{p\downarrow}^\dagger c_{p\uparrow}^\dagger  0\rangle$
0	0	$-t_{pd}$	$-t_{pd}$	$+t_{pd}$	$+t_{pd}$	0	0	$2(U_d + \Delta_{pd})$	$c_{2\downarrow}^\dagger c_{2\uparrow}^\dagger c_{1\downarrow}^\dagger c_{1\uparrow}^\dagger  0\rangle$



# superexchange: opposite spin



$$\begin{aligned}
 H_{\text{eff}} &= H_{00} + T_{01} \left( \varepsilon - \left( H_{11} + T_{12} (\varepsilon - H_{22})^{-1} T_{21} \right) \right)^{-1} T_{10} \\
 &\approx H_{00} - T_{01} H_{11}^{-1} T_{10} - T_{01} H_{11}^{-1} T_{12} H_{22}^{-1} T_{21} H_{11}^{-1} T_{10} \quad \text{expand in } 1/U_d \\
 &= -\frac{2t_{pd}^2}{U_d + \Delta_{pd}} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - \frac{2t_{pd}^4}{(U_d + \Delta_{pd})^2} \left( \frac{1}{U_d} + \frac{1}{U_d + \Delta_{pd}} \right) \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}
 \end{aligned}$$




singlet-triplet splitting:

$$J = \frac{4t_{pd}^4}{(U_d + \Delta_{pd})^2} \left( \frac{1}{U_d} + \frac{1}{U_d + \Delta_{pd}} \right)$$




# ferromagnetic superexchange



180° superexchange

hopping only via oxygen- $p$  pointing in direction connecting  $d$ -orbitals



90° superexchange

no hopping connecting  $d$ -orbitals but Coulomb exchange on oxygen


**double exchange**

# ferro superexchange: same spin



$$\begin{pmatrix} 0 & t_{pd} & t_{pd} & 0 \\ t_{pd} & U_d + \Delta_{pd} & 0 & t_{pd} \\ t_{pd} & 0 & U_d + \Delta_{pd} & t_{pd} \\ 0 & t_{pd} & t_{pd} & 2(U_d + \Delta_{pd}) - J_{xy} \end{pmatrix}$$

$$\begin{aligned} & c_{1\uparrow}^\dagger c_{x\downarrow}^\dagger c_{x\uparrow}^\dagger c_{y\downarrow}^\dagger c_{y\uparrow}^\dagger c_{2\uparrow}^\dagger |0\rangle \\ & c_{1\downarrow}^\dagger c_{1\uparrow}^\dagger c_{x\uparrow}^\dagger c_{y\downarrow}^\dagger c_{y\uparrow}^\dagger c_{2\uparrow}^\dagger |0\rangle \\ & c_{1\uparrow}^\dagger c_{x\downarrow}^\dagger c_{x\uparrow}^\dagger c_{y\uparrow}^\dagger c_{2\downarrow}^\dagger c_{2\uparrow}^\dagger |0\rangle \\ & c_{1\downarrow}^\dagger c_{1\uparrow}^\dagger c_{x\uparrow}^\dagger c_{y\uparrow}^\dagger c_{2\downarrow}^\dagger c_{2\downarrow}^\dagger |0\rangle \end{aligned}$$



$$H_{\text{eff}} = -\frac{2t_{pd}^2}{U_d + \Delta_{pd}} - \frac{4t_{pd}^4}{(U_d + \Delta_{pd})^2} \frac{1}{2(U_d + \Delta_{pd}) - J_{xy}}$$

# ferro superexchange: opposite spin $\uparrow\downarrow$ $\downarrow\uparrow$

$$\begin{pmatrix}
 0 & 0 & t_{pd} & 0 & t_{pd} & 0 & 0 & 0 \\
 0 & 0 & 0 & t_{pd} & 0 & t_{pd} & 0 & 0 \\
 t_{pd} & 0 & U_d + \Delta_{pd} & 0 & 0 & 0 & t_{pd} & 0 \\
 0 & t_{pd} & 0 & U_d + \Delta_{pd} & 0 & 0 & 0 & t_{pd} \\
 t_{pd} & 0 & 0 & 0 & U_d + \Delta_{pd} & 0 & t_{pd} & 0 \\
 0 & t_{pd} & 0 & 0 & 0 & U_d + \Delta_{pd} & 0 & t_{pd} \\
 0 & 0 & t_{pd} & 0 & t_{pd} & 0 & 2(U_d + \Delta_{pd}) - J_{xy} & \\
 0 & 0 & 0 & t_{pd} & 0 & t_{pd} & -J_{xy} & 2(U_d + \Delta_{pd})
 \end{pmatrix}
 \begin{matrix}
 c_{1\uparrow}^\dagger c_{x\downarrow}^\dagger c_{x\uparrow}^\dagger c_{y\downarrow}^\dagger c_{y\uparrow}^\dagger c_{2\downarrow}^\dagger |0\rangle \\
 c_{1\downarrow}^\dagger c_{x\downarrow}^\dagger c_{x\uparrow}^\dagger c_{y\downarrow}^\dagger c_{y\uparrow}^\dagger c_{2\uparrow}^\dagger |0\rangle \\
 c_{1\downarrow}^\dagger c_{1\uparrow}^\dagger c_{x\uparrow}^\dagger c_{y\downarrow}^\dagger c_{y\uparrow}^\dagger c_{2\downarrow}^\dagger |0\rangle \\
 c_{1\downarrow}^\dagger c_{1\uparrow}^\dagger c_{x\downarrow}^\dagger c_{y\downarrow}^\dagger c_{y\uparrow}^\dagger c_{2\uparrow}^\dagger |0\rangle \\
 c_{1\uparrow}^\dagger c_{x\downarrow}^\dagger c_{x\uparrow}^\dagger c_{y\downarrow}^\dagger c_{2\downarrow}^\dagger c_{2\uparrow}^\dagger |0\rangle \\
 c_{1\downarrow}^\dagger c_{x\downarrow}^\dagger c_{x\uparrow}^\dagger c_{y\uparrow}^\dagger c_{2\downarrow}^\dagger c_{2\uparrow}^\dagger |0\rangle \\
 c_{1\downarrow}^\dagger c_{1\uparrow}^\dagger c_{x\uparrow}^\dagger c_{y\downarrow}^\dagger c_{2\downarrow}^\dagger c_{2\uparrow}^\dagger |0\rangle \\
 c_{1\downarrow}^\dagger c_{1\uparrow}^\dagger c_{x\downarrow}^\dagger c_{y\uparrow}^\dagger c_{2\downarrow}^\dagger c_{2\uparrow}^\dagger |0\rangle
 \end{matrix}$$

$$\begin{aligned}
 H_{\text{eff}} &= -\frac{2t_{pd}^2}{U_d + \Delta_{pd}} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - \frac{4t_{pd}^4}{(U_d + \Delta_{pd})^2} \frac{1}{4(U_d + \Delta_{pd})^2 - J_{xy}^2} \begin{pmatrix} 2(U_d + \Delta_{pd}) & +J_{xy} \\ +J_{xy} & 2(U_d + \Delta_{pd}) \end{pmatrix} \\
 &= -\left( \frac{2t_{pd}^2}{U_d + \Delta_{pd}} + \frac{4t_{pd}^4}{(U_d + \Delta_{pd})^2} \frac{1}{2(U_d + \Delta_{pd}) - J_{xy}} \right) \text{ (as for same spin)} \\
 &\quad + \frac{4t_{pd}^4}{(U_d + \Delta_{pd})^2} \frac{J_{xy}}{4(U_d + \Delta_{pd})^2 - J_{xy}^2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}
 \end{aligned}$$

singlet-triplet splitting  $J = -\frac{4t_{pd}^4}{(U_d + \Delta_{pd})^2} \frac{2J_{xy}}{4(U_d + \Delta_{pd})^2 - J_{xy}^2}$


# double exchange

---


double exchange involves both, full Coulomb matrix and hopping

mixed-valence compound: non-integer filling of  $d$ -orbital  
 $d$ -electrons can hop even when  $U$  is large

simple model: two sites with two orbitals each



# double exchange




$$H = \begin{pmatrix} -J_{ab} & -t_{bb} \\ -t_{bb} & -J_{ab} \end{pmatrix}$$

$$\varepsilon_{\pm} = -J_{ab} \pm t_{bb}$$

$$\psi_{\pm} = \frac{1}{\sqrt{2}} \left( |\uparrow, \uparrow\rangle_1 |\cdot, \uparrow\rangle_2 \pm |\cdot, \uparrow\rangle_1 |\uparrow, \uparrow\rangle_2 \right) = \frac{1}{\sqrt{2}} \left( |\uparrow, \cdot\rangle_b \pm |\cdot, \uparrow\rangle_b \right) |\uparrow, \uparrow\rangle_a$$

*b*-electron hops against background of half-filled *a*-orbitals

# double exchange



$$H = \begin{pmatrix} -J_{ab} & -t_{bb} & 0 & 0 & 0 & 0 \\ -t_{bb} & 0 & -J_{ab} & 0 & 0 & 0 \\ 0 & -J_{ab} & 0 & -t_{bb} & 0 & 0 \\ 0 & 0 & -t_{bb} & 0 & -J_{ab} & 0 \\ 0 & 0 & 0 & -J_{ab} & 0 & -t_{bb} \\ 0 & 0 & 0 & 0 & -t_{bb} & -J_{ab} \end{pmatrix}$$

ground state  $\epsilon_0 = -J_{ab} - t_{bb}$


$$\frac{1}{\sqrt{6}} \left( |\uparrow, \uparrow\rangle_1 |\cdot, \downarrow\rangle_2 + |\cdot, \uparrow\rangle_1 |\uparrow, \downarrow\rangle_2 + |\cdot, \uparrow\rangle_1 |\downarrow, \uparrow\rangle_2 + |\downarrow, \uparrow\rangle_1 |\cdot, \uparrow\rangle_2 + |\uparrow, \downarrow\rangle_1 |\cdot, \uparrow\rangle_2 + |\cdot, \downarrow\rangle_1 |\uparrow, \uparrow\rangle_2 \right)$$

$$= \frac{1}{\sqrt{2}} \left( |\uparrow, \cdot\rangle_b + |\cdot, \uparrow\rangle_b \right) \frac{1}{\sqrt{2}} \left( |\uparrow, \downarrow\rangle_a + |\downarrow, \uparrow\rangle_a \right) + \frac{1}{\sqrt{2}} \left( |\downarrow, \cdot\rangle_b + |\cdot, \downarrow\rangle_b \right) |\uparrow, \uparrow\rangle_a$$


hopping electron aligns *a*-electrons ferromagnetically  
(teleports local triplet into triplet of *a*-electrons)

# double exchange


$S_z=3/2$




$S_z=1/2$



$S_z=-1/2$



$S_z=-3/2$



# double exchange

---

alternative model:

assume passive orbitals with many electrons (large Hund's rule spin)

example:  $e_g$  electrons hopping against  $t_{2g}$  background

consider these spins fixed with quantization axis tilted by  $\vartheta$  relative to each other



rotation of quantization axis

$$d_{2b\uparrow} = \cos(\vartheta/2) c_{2b\uparrow} - \sin(\vartheta/2) c_{2b\downarrow}$$

$$d_{2b\downarrow} = \sin(\vartheta/2) c_{2b\uparrow} + \cos(\vartheta/2) c_{2b\downarrow}$$


hopping mixes spins

$$-t_{bb} c_{2b\uparrow}^\dagger c_{1b\uparrow} = -t_{bb} \left( +\cos(\vartheta/2) d_{2b\uparrow}^\dagger + \sin(\vartheta/2) d_{2b\downarrow}^\dagger \right) c_{1b\uparrow}$$

$$-t_{bb} c_{2b\downarrow}^\dagger c_{1b\downarrow} = -t_{bb} \left( -\sin(\vartheta/2) d_{2b\uparrow}^\dagger + \cos(\vartheta/2) d_{2b\downarrow}^\dagger \right) c_{1b\downarrow}$$



# double exchange



assume  $a$ -spins cannot be flipped  $\Rightarrow$  no  $J$  terms

4 independent  $4 \times 4$  Hamiltonians


for  $t_{bb} \ll J_{ab}$  tilt merely reduces width of  $b$ -band

$$\varepsilon_{\pm} = -J_{ab} \pm t_{bb} \cos(\vartheta/2)$$

again, hopping of  $b$ -electron prefers ferro-aligned  $a$ -electrons

# orbital ordering

same model, but now one electron per orbital




$$H = \left( \begin{array}{cc|cc} 0 & 0 & -t_{bb} & -t_{aa} \\ 0 & 0 & +t_{aa} & +t_{bb} \\ \hline -t_{bb} & +t_{aa} & U_{ab} - J_{ab} & 0 \\ -t_{aa} & +t_{bb} & 0 & U_{ab} - J_{ab} \end{array} \right)$$

$$H_{\text{eff}} \approx -\frac{1}{U_{ab} - J_{ab}} \begin{pmatrix} t_{aa}^2 + t_{bb}^2 & -2t_{aa}t_{bb} \\ -2t_{aa}t_{bb} & t_{aa}^2 + t_{bb}^2 \end{pmatrix} = -\frac{(t_{aa} - t_{bb})^2}{U_{ab} - J_{ab}} - \frac{2t_{aa}t_{bb}}{U_{ab} - J_{ab}} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}$$

effective interaction between orbitals: orbital singlet/triplet

# orbital ordering: opposite spins



$$H = \left( \begin{array}{cccc|cccc} 0 & 0 & 0 & 0 & -t_{bb} & -t_{aa} & 0 & 0 \\ 0 & 0 & 0 & 0 & +t_{aa} & +t_{bb} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -t_{bb} & -t_{aa} \\ 0 & 0 & 0 & 0 & 0 & 0 & +t_{aa} & +t_{bb} \\ \hline -t_{bb} & +t_{aa} & 0 & 0 & U_{ab} & 0 & -J_{ab} & 0 \\ -t_{aa} & +t_{bb} & 0 & 0 & 0 & U_{ab} & 0 & -J_{ab} \\ 0 & 0 & -t_{bb} & +t_{aa} & -J_{ab} & 0 & U_{ab} & 0 \\ 0 & 0 & -t_{aa} & +t_{bb} & 0 & -J_{ab} & 0 & U_{ab} \end{array} \right)$$


# orbital-ordering: opposite spin

$$\begin{aligned}
 H_{\text{eff}} &\approx -\frac{1}{U_{ab}^2 - J_{ab}^2} \begin{pmatrix} (t_{aa}^2 + t_{bb}^2)U_{ab} & -2t_{aa}t_{bb}U_{ab} & (t_{aa}^2 + t_{bb}^2)J_{ab} & -2t_{aa}t_{bb}J_{ab} \\ -2t_{aa}t_{bb}U_{ab} & (t_{aa}^2 + t_{bb}^2)U_{ab} & -2t_{aa}t_{bb}J_{ab} & (t_{aa}^2 + t_{bb}^2)J_{ab} \\ (t_{aa}^2 + t_{bb}^2)J_{ab} & -2t_{aa}t_{bb}J_{ab} & (t_{aa}^2 + t_{bb}^2)U_{ab} & -2t_{aa}t_{bb}U_{ab} \\ -2t_{aa}t_{bb}J_{ab} & (t_{aa}^2 + t_{bb}^2)J_{ab} & -2t_{aa}t_{bb}U_{ab} & (t_{aa}^2 + t_{bb}^2)J_{ab} \end{pmatrix} \\
 &= -\frac{1}{U_{ab}^2 - J_{ab}^2} \begin{pmatrix} U_{ab} & J_{ab} \\ J_{ab} & U_{ab} \end{pmatrix} \otimes \begin{pmatrix} t_{aa}^2 + t_{bb}^2 & -2t_{aa}t_{bb} \\ -2t_{aa}t_{bb} & t_{aa}^2 + t_{bb}^2 \end{pmatrix} \\
 &= -\frac{1}{U_{ab}^2 - J_{ab}^2} \left[ U_{ab} + J_{ab} - J_{ab} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \right] \otimes \left[ (t_{aa} - t_{bb})^2 + 2t_{aa}t_{bb} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \right]
 \end{aligned}$$

spin-exchange

orbital-exchange

simultaneous coupling of spins and orbital occupations  
 spin- and orbital-exchange tend to have opposite sign



# summary

---

## exchange mechanisms

dominant magnetic interaction in materials

not a fundamental but an **effective interaction**: model/mechanism

- ▶ Coulomb exchange: off-diagonal Coulomb matrix-elements;  
ferromagnetic coupling (Hund's rule)
- ▶ kinetic exchange: only diagonal Coulomb matrix-elements & hopping
- ▶ direct exchange: anti-ferromagnetic spins: virtual hopping  $-4t^2/U$
- ▶ superexchange: hopping via O-*p* orbitals  
tends to be anti-ferromagnetic (180° superexchange)  
but 90° superexchange is ferromagnetic
- ▶ double exchange: hopping electrons align spins ferromagnetically
- ▶ orbital ordering: exchange interaction between orbital occupations


# summary

$$H_U = \begin{pmatrix} U_{ab} - J_{ab} & 0 & 0 & 0 \\ 0 & U_{ab} & -J_{ab} & 0 \\ 0 & -J_{ab} & U_{ab} & 0 \\ 0 & 0 & 0 & U_{ab} - J_{ab} \end{pmatrix}$$

direct exchange



Coulomb exchange:  
ferro (Hund's rule)

kinetic exchange:  
anti-ferro



# summary

double exchange: often ferro



orbital-ordering

