variational principle and Schrödinger equation

normalized wavefunctions

energy functional $E[\psi] = \langle \psi | H | \psi
angle$

variation

 $E[\psi + \delta \psi] = \langle \psi | H | \psi \rangle + \langle \delta \psi | H | \psi \rangle + \langle \psi | H | \delta \psi \rangle + \langle \delta \psi | H | \delta \psi \rangle$ = $E[\psi] + \delta E[\psi] + \mathcal{O}^2(\delta \phi)$

functional derivative

$$rac{\delta E[\psi]}{\delta \psi} = rac{E[\psi + \delta \psi] - E[\psi]}{\delta \psi}$$

ensure normalization via Lagrange multiplier

$$L[\psi] = E[\psi] - \varepsilon \left(\langle \psi | \psi \rangle - 1 \right)$$

variational principle $0 = \delta L[\psi] = \langle \delta \psi | \left(H | \psi \rangle - \varepsilon | \psi \rangle \right) + \left(\langle \psi | H - \varepsilon \langle \psi | \right) | \delta \psi \rangle$

gives time-independent Schrödinger equation

variational principle and finite basis set

trial function
$$|\psi\rangle = \sum_{n} \alpha_{n} |\psi_{n}\rangle$$
 (orthogonal basis)
 $E[\psi] = \sum_{n,m} \alpha_{n} \alpha_{m} \langle \psi_{n} | H | \psi_{m} \rangle$
 $L[\psi] = \sum_{n,m} \alpha_{n} H_{nm} \alpha_{m} - \varepsilon \left(\sum \alpha_{n}^{2} - 1\right)$

variational principle

$$0 = \frac{\partial L}{\partial \alpha_{k}} = \sum_{n} \alpha_{n} H_{nk} + \sum_{m} H_{km} \alpha_{m} - 2\varepsilon \alpha_{k} = \left(\sum_{n} \alpha_{n} H_{nk} - \varepsilon \alpha_{k}\right) + \left(\sum_{m} H_{km} \alpha_{m} - \varepsilon \alpha_{k}\right)$$

$$\begin{pmatrix} H_{11} & H_{12} & \cdots & H_{1N} \\ H_{21} & H_{22} & \cdots & H_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ H_{N1} & H_{N2} & \cdots & H_{NN} \end{pmatrix} \begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \vdots \\ \alpha_N \end{pmatrix} = \varepsilon \begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \vdots \\ \alpha_N \end{pmatrix}$$

non-orthogonal basis set

trial function
$$|\psi\rangle = \sum_{n} \alpha_{n} |\psi_{n}\rangle$$
 (arbitrary basis)
overlap matrix $S_{nm} = \langle \psi_{n} | \psi_{m} \rangle$
 $L[\psi] = \sum \alpha_{n} H_{nm} \alpha_{m} - \varepsilon \left(\sum \alpha_{n} S_{nm} \alpha_{m} - 1 \right)$

variation results in generalized eigenvalue problem

$$\begin{pmatrix} H_{11} & H_{12} & \cdots & H_{1N} \\ H_{21} & H_{22} & \cdots & H_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ H_{N1} & H_{N2} & \cdots & H_{NN} \end{pmatrix} \begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \vdots \\ \alpha_N \end{pmatrix} = \varepsilon \begin{pmatrix} S_{11} & S_{12} & \cdots & S_{1N} \\ S_{21} & S_{22} & \cdots & S_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ S_{N1} & S_{N2} & \cdots & S_{NN} \end{pmatrix} \begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \vdots \\ \alpha_N \end{pmatrix}$$

tight-binding

$$H = -\frac{1}{2}\Delta + \sum V_n$$

superposition of well separated potentials V_n

use solutions of individual potentials

$$H_n = -rac{1}{2}\Delta + V_n \qquad H_n |n
angle = arepsilon_n |n
angle$$
as basis set

tight-binding approximation for matrix elements:

$$\langle n|H|n \rangle = \varepsilon_n + \sum_{m \neq n} \langle n|V_m|n \rangle$$

$$\langle n|H|m \rangle = (\varepsilon_n + \varepsilon_m)/2 \langle n|m \rangle + \langle n|V_n + V_m|m \rangle/2 + \sum_{k \neq n,m} \langle n|V_k|m \rangle$$

for *n* and *m* nearest neighbors

Hellmann-Feynman theorem & forces

molecular dynamics:

 $H(R_{\alpha})|\Psi(R_{\alpha})\rangle = E(R_{\alpha})|\Psi(R_{\alpha})\rangle$

force on nucleus α $\vec{F}_{\alpha} = -\vec{\nabla}E(R_{\alpha}) = -\langle \Psi(R_{\alpha}) | \vec{\nabla}H | \Psi(R_{\alpha}) \rangle$



for non-degenerate state ...

Hellmann-Feynman theorem

degenerate perturbation theory in Phys. Rev. B 2003 !

PHYSICAL REVIEW B 68, 033105 (2003)

Hellmann-Feynman theorem at degeneracies

Ofir E. Alon and L. S. Cederbaum

Theoretische Chemie, Physikalisch-Chemisches Institut, Universität Heidelberg, Im Neuenheimer Feld 229, D-69120 Heidelberg, Germany (Received 2 August 2002; revised manuscript received 18 March 2003; published 30 July 2003)

The Hellmann-Feynman theorem is extended to account for degenerate states. Given a point $\lambda = \lambda_0$ in parameter space where the energy $E(\lambda_0)$ is *n*-fold degenerate, it is shown that the corresponding *n* forces (slopes) are obtained by diagonalizing the derivative of the Hamiltonian, $-\partial H(\lambda)/\partial \lambda|_{\lambda=\lambda_0}$, in the subspace of degenerate eigenstates. Such a rotation within the subspace of degenerate eigenfunctions is easy and simple to apply in practical calculations and should be performed *separately* for each independent direction in parameter space for which the forces are to be calculated.

DOI: 10.1103/PhysRevB.68.033105

Hellmann-Feynman theorem $(HFT)^{1,2}$ is widely used in practice to calculate forces (slopes) in atoms, molecules and crystals, but the way it appears in Refs. 1 and 2 and in quantum mechanics textbooks (see, e.g., Ref. 3) is only correct for nondegenerate states (see below). It should be mentioned that there exists a more general form of HFT, which is defined also for off-diagonal matrix elements (see, e.g., Refs. 4-9). However, the prescribed expressions do not provide an PACS number(s): 31.15.Md, 03.65.-w, 71.15.-m

depend also on the *direction* (in parameter space) for which one calculates the forces. Practically, a set of degenerate eigenfunctions obtained in a numerical calculation has to be appropriately rotated to allow for the correct calculation of forces as expectation values of $-\partial H(\lambda)/\partial \lambda$ [see Eq. (1)] at degeneracy. These detailed and relevant points, all related to the "freedom" to chose the linear combinations of eigenstates at degeneracy, have been overlooked previously.

Hellmann-Feynman theorem II

a surprising failure of reason...

PHYSICAL REVIEW B 66, 033110 (2002)

Breakdown of the Hellmann-Feynman theorem: Degeneracy is the key

G. P. Zhang

Department of Physics, Indiana State University, Terre Haute, Indiana 47809

Thomas F. George

Office of the Chancellor/Departments of Chemistry and Physics & Astronomy, University of Wisconsin-Stevens Point, Stevens Point, Wisconsin 54481-3897

(Received 17 May 2002; published 29 July 2002)

The Hellmann-Feynman theorem is a powerful and popular method to efficiently calculate forces in a variety of dynamical processes, but its validity has rarely been addressed. Here a surprising failure of this theorem is reported. The forces calculated by the theorem can be more than fifty times smaller than the forces calculated by the finite differential method. Numerical evidence shows that the energy-level degeneracy is the main reason. An analytical proof reveals that although eigenvalues do not depend on a linear combination of degenerate wave functions, forces do sensitively depend on it, which leads to ill-defined forces. A scheme is proposed to overcome this difficulty.

DOI: 10.1103/PhysRevB.66.033110

The Hellmann-Feynman theorem¹ is powerful and has been widely used in many fields such as dynamical processes,² molecular dynamics,³ chemical reactions, and PACS number(s): 71.15.-m

merical examples for C_{60} explicitly show that forces calculated by the Hellmann-Feynman theorem can be fifty times smaller than forces calculated by the finite differential method itself also gives

Hellmann-Feynman theorem III

an elegant method...

PHYSICAL REVIEW B 69, 167102 (2004)

Extended Hellmann-Feynman theorem for degenerate eigenstates

G. P. Zhang

Department of Physics, Indiana State University, Terre Haute, Indiana 47809, USA

Thomas F. George

Office of the Chancellor/Departments of Chemistry & Biochemistry and Physics & Astronomy, University of Missouri-St. Louis, St. Louis, Missouri 63121, USA

(Received 2 August 2003; revised manuscript received 26 January 2004; published 27 April 2004)

In a previous paper, we reported a failure of the traditional Hellmann-Feynman theorem (HFT) for degenerate eigenstates. This has generated enormous interest among different groups. In four independent papers by Fernandez, by Balawender, Hola, and March, by Vatsya, and by Alon and Cederbaum, an elegant method to solve the problem was devised. The main idea is that one has to construct and diagonalize the force matrix for the degenerate case, and only the eigenforces are well defined. We believe this is an important extension to HFT. Using our previous example for an energy level of fivefold degeneracy, we find that those eigenforces correctly reflect the symmetry of the molecule.

DOI: 10.1103/PhysRevB.69.167102

The Hellmann-Feynman theorem (HFT) (Ref. 1) is an efficient method, but when eigenstates are degenerate, an extension is necessary. The reason is that any good linear comPACS number(s): 71.15.-m

culated from two different configurations of C_{60} , the pristine and excited C_{60} . The fivefold degenerate case is from the pristine C_{60} . We present the force matrix elements in Table I.