

### Exercise Sheet 8

#### 1. Spherical potential approximation

- i. Show that the charge density of a filled shell  $n(\vec{r}) = \sum_{m=-l}^l |\varphi_{nlm}(\vec{r})|^2$  is spherically symmetric.
- ii. Calculate the spherical average of the charge density of an open shell with  $N_{occ}$  electrons with fixed quantum numbers  $(n, l)$  but arbitrary  $m$

$$n(r) := \frac{1}{4\pi} \int_0^{2\pi} d\varphi \int_0^\pi \sin\vartheta d\vartheta n(\vec{r}).$$

in terms of the radial function  $u_{nl}(r)$ .

#### 2. Slater-Condon parameters

Calculate the Slater-Condon parameters

$$F_{nl}^{(k)} = \int_0^\infty dr_1 u_{nl}^2(r_1) \left( \frac{1}{r_1^{k+1}} \int_0^{r_1} dr_2 u_{nl}^2(r_2) r_2^k + r_1^k \int_{r_1}^\infty dr_2 u_{nl}^2(r_2) \frac{1}{r_2^{k+1}} \right)$$

for the 3s (only  $k = 0$ ), 3p ( $k = 0, 2$ ) and 3d ( $k = 0, 2, 4$ ) functions of hydrogen. Give the resulting energies in eV. For a given shell  $(n, l)$ , calculate the ratio of the Parameters  $F^{(k')}/F^{(k)}$ .